


**WEST DESERT  
GREATER SAGE-GROUSE  
(CENTROCERCUS UROPHASIANUS)  
LOCAL CONSERVATION PLAN**



**July, 2007**

**West Desert Adaptive Resource Management Local Working Group**

**WEST DESERT  
GREATER SAGE-GROUSE  
(*CENTROCERCUS UROPHASIANUS*)  
LOCAL CONSERVATION PLAN**

**May 7, 2007**

**West Desert Adaptive Resource Management Local Working Group**

*Published by*

**Utah State University Extension  
UMC 5230  
Logan, Utah 84322-5230**

*and*

**Jack H. Berryman Institute  
UMC 5230  
Logan, Utah 84322-5230**

*and*

**Utah Division of Wildlife Resources  
1594 W. North Temple  
Salt Lake City, Utah 84116**

This report should be cited as:

West Desert Adaptive Resource Management Local Working Group (WDARM). 2007. West Desert Greater Sage-grouse (*Centrocercus urophasianus*) Local Conservation Plan. Utah State University Extension and Jack H. Berryman Institute and Utah Division of Wildlife Resources. Salt Lake City, Utah. Unpublished Report.

**West Desert Greater Sage-grouse (*Centrocercus urophasianus*)  
Local Conservation Plan**

**Table of Contents**

	<b>Page</b>
List of Figures .....	i
List of Tables .....	ii
I. Executive Summary.....	1
II. Introduction.....	2
A. Purpose .....	2
B. Goals and Scope.....	3
C. Plan Duration .....	4
D. West Desert Adaptive Resource Management Local Working Group.....	4
E. Socioeconomic Considerations Including Consequences of Federal Listing .....	6
F. Management and Legal Authorities .....	7
Utah Division of Wildlife Resources (UDWR) .....	7
Counties.....	8
Natural Resources Conservation Service (NRCS).....	8
Bureau of Land Management (BLM).....	9
School and Institutional Trust Lands Administration (SITLA) .....	9
United States Forest Service (USFS) .....	10
Memorandum of Understanding .....	10
G. Policy for Evaluation of Conservation (PECE) Standards .....	11
III. Conservation Assessment .....	12
A. General Sage-grouse Biology/Ecology.....	12
C. Distribution and Abundance .....	17
Historic Distribution of Sage-grouse.....	17
Lek Counts .....	17
D. Assessment of Local Population.....	19
Plan Area .....	19
Population Status and Distribution.....	23
Local Habitat .....	25
Habitat Improvements & Completed Conservation Actions.....	25
IV. Threat Analysis.....	31
A. Altered Water Distribution .....	31
B. Development and Human Infrastructure.....	32
C. Drought and Weather .....	35
D. Hunting .....	37
E. Fire .....	40
F. Livestock Grazing .....	41
G. OHV Recreation .....	44
H. Invasive/Noxious Weeds .....	46
I. Parasitism & Disease.....	47
J. Predation .....	49
K. Renewable and Non-renewable Energy Development.....	59
L. Vegetation Management .....	61
V. Conservation Strategy .....	62
A. Strategies and Actions .....	62
B. Priority Evaluation .....	67
IV. Literature Cited.....	69

Appendix A..... 81  
Appendix B..... 82

**List of Figures**

Figure 1. The WDARM Resource Area with land management and landownership designated.20  
Figure 2. Maximum total number of males counted on all leks in the Vernon Subunit of the Resource Area, 1968-2006..... 24  
Figure 4. Location of sage-grouse nesting and brood-rearing habitat in the WDARM Resource Area, as identified by the UDWR, 2006. .... 26  
Figure 5. Location of sage-grouse winter habitat in the WDARM Resource Area, as identified by the UDWR, 2006. .... 27  
Figure 6. Location of Utah Big Game Range Trend Sites and vegetation monitoring sites located within sage-grouse seasonal habitat, as identified by the UDWR, 2006. .... 28  
Figure 8. Average precipitation for the Weber/Ogden River Basin 1995-2005 and the 30-year average (Utah Division of Water Resources 2006). .... 36  
Figure 9. USDA-WS reported coyote take in Utah, 1917-2004. Data reported by USDA-WS. 51  
Figure 10. USDA-WS reported fox take in Utah, 1972-2004. Data reported by USDA-WS. ... 52  
Figure 11. USDA-WS reported skunk take in Utah, 1917-2004. Data reported by USDA-WS.54  
Figure 13. Location of drill sites in the WDARM Resource Area. Data obtained from [http://www.ogm.utah.gov/oilgas/MAP\\_SEARCH/map\\_search.htm](http://www.ogm.utah.gov/oilgas/MAP_SEARCH/map_search.htm)..... 60

**List of Tables**

Table 1. West Desert Adaptive Resource Management Local Working Group partners.....5

Table 2. Landownership in the WDARM Resource Area.....19

Table 3. Habitat improvement projects implemented in 2004 and 2005 and proposed for 2006 by the WDARM partners on public and private land in the Resource Area.....30

Table 4. Sage-grouse hunter harvest information for Tooele County, Utah 1984-2003 (UDWR 1984-2000, UDWR 2003). .....38

Table 5. Sage-grouse hunter harvest information for Juab County, Utah 1984-2003 (UDWR 1984-2000, UDWR 2003). .....39

Table 5. Relative importance/contribution of individual threats to reducing or degrading aspects of sage-grouse populations in the WDARM Resource Area. Threats are described in the “Threat Analysis” section of this Plan. Rankings are as follows: L = low; M = medium; H = high; and VH = very high. Ranks are defined according to TNC (2005). .....68

## **I. Executive Summary**

The West Desert Greater Sage-grouse Conservation Plan (Plan) is the culmination of a two-year effort by the West Desert Adaptive Resource Management Local Working Group (WDARM). WDARM members include representatives from state and federal land management and resource agencies, as well as private landowners. WDARM formed in 2005 to proactively manage Greater Sage-grouse (*Centrocercus urophasianus*) populations and their habitats in response to increasing concern about the status of sage-grouse populations rangewide and within their local area. The impetus for the writing of this Plan came from a mandate by the Utah Division of Wildlife Resources (UDWR) in their Statewide Strategic Management Plan, passed by the Wildlife Board in 2002.

The Plan will provide an assessment of the status of Tooele and Juab Counties sage-grouse population. The intent of the Plan is to provide guidance and recommendations to meet the overall goal of maintaining and, where possible, increasing sage-grouse populations and improving habitat conditions in Tooele and Juab Counties. The Plan is designed to meet the guidelines set forth by the United States Fish and Wildlife Service (USFWS) in their Policy for Evaluation of Conservation Efforts (PECE) standards.

The Plan directly and indirectly addresses the five USFWS listing factors as they apply to Greater Sage-grouse in Tooele and Juab Counties. Recommendations and guidance suggested within the Plan can be adopted by all WDARM partners on a voluntary basis. WDARM encourages participation and adoption of these practices, where applicable, by private landowners in the local area. Participation by private landowners and consideration of landowner needs are critical for management of sage-grouse populations and habitat located on private lands and will be instrumental in meeting the overall goals of the Plan. True success will only be achieved by managing on a landscape scale. The Plan provides an opportunity to promote ecologically sound management of private and public lands for sage-grouse without impinging on private property rights.

Information contained in the Plan is based on a thorough review of the published and unpublished literature relevant to sage-grouse and sagebrush habitats, and also on the intimate, local knowledge possessed by WDARM partners who live and work in the area. Because a wealth of general information exists about sage-grouse and is available in published documents (Connelly et al. 2000, Connelly et al. 2004), we provide only a brief overview of general sage-grouse ecology and try instead to focus on conditions and issues specific to Tooele and Juab Counties. Knowledge gaps are also identified.

WDARM analyzed threats currently or potentially affecting sage-grouse and sagebrush habitats in Tooele and Juab Counties. The Threat Analysis, combined with recommended strategies and actions, provides a framework for implementation of the Plan over the next ten years by WDARM partners. Implementation will be conducted within an adaptive resource management framework. As relevant information from a local and rangewide perspective becomes available, it will be used to modify and refine management strategies, priorities, and general understanding of sage-grouse ecology in the area. Annual evaluation and reporting performed by WDARM will track progress on the objectives outlined in this Plan.

## **II. Introduction**

### **A. Purpose**

The mission of the West Desert Adaptive Resource Management Sage-grouse Conservation Plan is to aid in reaching the goal of maintaining and improving current abundance and viability of Greater Sage-grouse (hereafter referred to as sage-grouse) populations and their habitat in Tooele and Juab Counties. Implementation of the Plan will take into consideration historic land uses and long-term social and economic issues. The Plan will help meet this goal by providing management solutions based on local or compatible data and research to the extent practical. In addition, WDARM hopes to develop management solutions that will result in diverse and productive sagebrush habitat for sage-grouse, while simultaneously recognizing that healthy sagebrush habitats are valuable to the existence of other species. The Plan will identify management areas, key local issues, conservation strategies, population information, research and monitoring needs, and support long-term funding. Adaptive management will be used to maintain the Plan as a continuously evolving document. The Plan will coordinate development of project proposals with the Central Region Utah Partners for Conservation and Development Regional Team to maintain and enhance sage-grouse habitat.

This Plan was called for in, and builds upon, the Utah Greater Sage-grouse Strategic Management Plan (Strategic Plan), passed by the Utah Wildlife Board in 2002. The Strategic Plan was developed by the Utah Greater Sage-grouse Working Group, which included representatives from state and federal natural resource agencies, and local conservation organizations concerned with the health and proper management of Greater Sage-grouse and sagebrush-steppe ecosystems throughout Utah. The primary purpose of the Strategic Plan was to address declining populations of sage-grouse and to develop a framework for agencies to work within. Further, the Strategic Plan identifies certain management units throughout the state where Adaptive Resource Management Local Working Groups could be organized to identify local issues and implement local adaptive resource management plans. These groups will work collaboratively to address declining sage-grouse populations, and the loss, degradation, and fragmentation of sagebrush steppe communities, with the end goal of protecting and conserving these and other natural resources.

The Plan is designed to meet the guidelines set forth by the USFWS in their PECE standards. The USFWS uses PECE standards as a guideline to evaluate whether conservation plans will be considered when making listing and listing priority decisions. The Plan was written to address the USFWS five Listing Factors which are:

1. Present or threatened destruction, modification, or curtailment of its habitat or range
2. Over-utilization for commercial, recreational, scientific, or educational purposes
3. Disease or predation
4. Authorities and inadequacy of existing regulatory mechanisms
5. Other natural or man-made factors affecting its continued existence

The Plan directly and indirectly addresses the five USFWS listing factors as they apply to sage-grouse in the West Desert. In addition, the Plan will identify issues, potential strategies, and provide for implementation of proposed conservation actions. The Plan is neither a National Environmental Policy Act (NEPA) decision document, nor a federal or state recovery plan. Any

Candidate Conservation Agreement with Assurances developed by the UDWR will be based on the Plan, but will include the NEPA process. Use of the Plan by agencies, private enterprise, and private individuals is strictly voluntary. State and federal resource management agencies involved with sage-grouse management, however, are required to manage sage-grouse populations and habitat by various state and federal statutes and policies. The information contained in the Plan is intended to serve as a set of guidelines for those state and federal agencies to maintain and enhance sage-grouse habitat and sage-grouse populations in the West Desert. Participation by private landowners and consideration of landowner needs are critical for management of sage-grouse populations and their habitat located on private lands and will be of great importance to meeting the overall goals of the Plan. True success will only be achieved by managing on a landscape scale. The Plan provides an opportunity to promote ecologically sound management of private and public lands for sage-grouse, without impinging on private property rights.

It is the intent of WDARM that this Plan be read and interpreted in its entirety. If the reader reads only isolated sections of this Plan, single statements may be taken out of context or misinterpreted.

## **B. Goals and Scope**

The goals of the Plan are separated into two categories: Assessment Goals and Strategy Goals. The goals are not listed in any particular order.

### Assessment Goals:

The Plan will provide an assessment of the status of the sage-grouse population in the West Desert by accomplishing the following goals:

1. Estimate current population size and evaluate population trends; estimate amount and condition of habitat.
2. Identify research needs and knowledge gaps.
3. Determine population and habitat needs for the future.
4. Identify and discuss threats that have potential to impact sage-grouse in the West Desert, especially those associated with the five USFWS Listing Factors.

### Strategy Goals:

The intent of the Plan is to maintain and, where possible, increase sage-grouse populations and improve habitat conditions in the West Desert by carrying out the following goals:

1. Incorporate management strategies from state and federal agency partners, local governments, and established rangewide conservation and management guidelines (Connelly et al. 2000, Connelly et al. 2004).
2. Increase effective communication with all potential stakeholders in the West Desert and the state of Utah, through outreach, information distribution, and education.
3. Address and prioritize threats to aid in prioritizing management solutions.
4. Identify and pursue funding sources, or support partners in their pursuance of funding for projects that will help achieve specific strategies and actions.

## Scope

This Plan is designed to span multiple land ownerships and multiple land uses throughout the defined geographic area. It is hoped that through implementation of this adaptive plan, specific conservation issues will be addressed, implemented, and monitored across geographic and political boundaries to increase consistency of the practices implemented and information collected. The assessment and strategies described herein are specific to Tooele and Juab Counties, and were developed with the unique ecological, social, and economic concerns of that area in mind. A detailed description of the West Desert Resource Area is provided later in the Plan.

## **C. Plan Duration**

The Plan was designed and written to be a dynamic document that can adapt with the needs of the local sage-grouse population, habitats, and local community as necessary. WDARM will re-evaluate sage-grouse populations and habitats, and will review progress on strategies listed in the Plan as per the Standard Operating Procedures (SOP; Appendix A). The Plan was written to support conservation actions over a ten-year period. Early termination of the Plan would occur if the sage-grouse was listed under the Endangered Species Act (ESA) or if sage-grouse were removed from the UDWR's Sensitive Species list. Species on the Sensitive Species list include species that are federally listed, are candidates for federal listing, or for which there is "credible scientific evidence to substantiate a threat to continued population viability" (Utah Division of Wildlife Resources 2005).

## **D. West Desert Adaptive Resource Management Local Working Group**

As a result of the Strategic Plan, the WDARM was formed in 2004 and has worked consistently and cooperatively toward the completion and implementation of the Plan since that time. WDARM was organized and facilitated by Scott Pratt of Utah State University (USU) Extension Services and later by Sarah G. Lupis of Utah's Community-Based Conservation Program (CBCP), a collaborative partnership between the UDWR and USU Extension Services with support from the Jack H. Berryman Institute. Ms. Lupis also served as the technical writer and compiler of the Plan itself. WDARM is comprised of state and federal agency personnel, representatives from local government, non-profit organizations, academic institutions, private industry, and private individuals. The agencies, organizations, and individuals who contributed to the Plan through their participation in WDARM are listed in Table 1. When 'we' or 'our' is used in the Plan, it refers to WDARM.

The role of WDARM participants is to guide the development of the Plan and to represent their agencies. After completion of the plan, WDARM participants will continue to meet to update the Plan, and through an adaptive process incorporate the results of research and monitoring efforts, new information, and lessons learned. Guidance for continued operation of WDARM can be found in the SOP (Appendix A). The director of the UDWR has the ultimate authority for the Plan.

Prior to writing the Plan, we reviewed several local sage-grouse conservation plans, statewide plans, and rangewide plans and assessments (UDWR 2002, Armentrout et al. 2004, Lincoln

*West Desert Greater Sage-grouse Conservation Plan*

County Sage-grouse Technical Review Team 2004, Northwest Colorado Greater Sage-Grouse Working Group 2004), to determine the most appropriate structure and content. In addition, a thorough literature review was conducted to ensure that the Plan contained the most recent information available on sage-grouse ecology, life history, and habitat requirements. Annual working group meetings, work plans, and accomplishment reports will monitor progress toward meeting the goals of the Plan. The Plan is intended to be an evolving document. Incorporating principles of adaptive management and changing as new information arises, will help to ensure success of the Plan and WDARM.

Table 1. West Desert Adaptive Resource Management Local Working Group partners.

Bureau of Land Management (BLM)
Farm Bureau Federation
Farm Services Agency (FSA)
Tooele County Commission
Natural Resource Conservation Service (NRCS)
Juab County Commission
Bennion Ranch
The Nature Conservancy (TNC)
USDA Forest Service (USFS)
USDA Wildlife Services (WS)
U.S. Fish and Wildlife Service (USFWS)
Utah Division of Wildlife Resources (UDWR)
Utah School and Institutional Trust Lands Administration (SITLA)
Utah State University Extension (USU/EXT)
Deep Creek Watershed Council
Tooele County Natural Resources Local Working Group
Shambip Soil Conservation District
Utah Association of Conservation Districts
Goshute Tribe

Management strategies and recommendations described in the Plan will be periodically updated to incorporate results of research efforts, new information, and the results of management actions through annual reviews and progress reports.

WDARM operates through an open public process based on consensus decision making. For decisions regarding the Plan, consensus was reached by participating members and/or those present at the time the decision was made. Sections 5 and 6 of the ESA, direct state and federal agencies to cooperate in developing conservation activities that protect candidate species. Because the responsibility lies with state and federal agencies, ultimately decisions are limited to them. However, all agencies felt that it was important to involve the public in the decision-making and planning process to the greatest extent possible. The importance of public-private partnerships was highlighted in the Statewide Strategic Management Plan (UDWR 2002):

“An important part of solving the habitat management problems that face sage-grouse, is to work together closely so that all landowners and land managers are aware of the needs of local populations and how to meet them.”

WDARM provided regular opportunities for public involvement, participation, and comment on the Plan. Regular meetings were scheduled to meet the needs of the greatest number of WDARM participants possible. Meetings were announced by direct mailings, on the CBCP website ([www.extension.usu.edu/cbcp](http://www.extension.usu.edu/cbcp)), via email, and through personal phone calls and invitations. During the planning process, WDARM met at least every other month and often monthly. Meeting minutes and critical updates were provided via email, direct mailing, and on the CBCP website. In addition, an annual community forum was held to update the local community on WDARM’s activities and solicit participation and comment from local stakeholders. Annual forums were announced in a similar fashion as regular meetings. The CBCP provided informational material to County Extension offices for display and distribution to the local community, and CBCP personnel met regularly with County governments (commissions and councils) to update them on WDARM’s activities and Plan progress. The final draft of the Plan was made available to all potential stakeholders that WDARM was aware of, and comments were encouraged.

#### **E. Socioeconomic Considerations Including Consequences of Federal Listing**

Communities in the Intermountain West are reflective of diverse and complicated relationships between natural resource extraction industries (agriculture, minerals, energy development, etc.), landownership (private vs. public), and local, state, and federal laws and regulations. These rural communities are reflective of cyclic (boom/bust) economies and global economics that drive commodity prices. To be successful, management recommendations and solutions designed to improve sage-grouse populations and habitats must reflect and be sensitive to local socioeconomic issues.

State and federal agencies will coordinate with local landowners, county, and local governments to develop solutions that will meet ecological requirements while maintaining social and economic values of the local community to the greatest extent possible. Participation by local stakeholders in the planning process has helped to ensure that recommendations and guidelines presented in the Plan will meet the needs of the local community. In many instances, cooperation between landowners and agencies results in more useful and cost-effective habitat improvement projects that ultimately benefit both sage-grouse and local interests.

Listing the sage-grouse under the provisions of ESA could have a variety of local impacts. Activities potentially affected include noxious weed control, maintenance of rights of way, subdivisions and land development, livestock grazing management, big game wildlife management, and recreational land use. Broadly applying ‘take’ regulations under ESA could have a significant local impact. There will likely be an increase in bureaucratic processes in environmental permitting and compliance. Ultimately, the listing could result in slowed growth and the elimination of new projects because of the increased cost of environmental permitting and compliance.

In the event of listing, this Plan, along with other local conservation plans, statewide

conservation plans, and rangewide conservation assessments and strategies will be used by the USFWS to develop a federal recovery plan. Should these events transpire, the USFWS would also strive to consider social and economic needs to the maximum extent possible. In the July 1, 1994 Federal Register (59 FR 34272), the USFWS issued a policy to involve stakeholders in the preparation of federal recovery plans to help minimize the social and economic impacts of implementing recovery actions.

## **F. Management and Legal Authorities**

Existing state, federal, and county regulations offer protection to sage-grouse in the Uintah Basin. State laws restrict possession of individual birds. Funding programs in Utah support population and habitat conservation and monitoring activities. Federal agencies including the Bureau of Land Management (BLM), U.S. Forest Service (USFS), National Park Service (NPS), Natural Resources Conservation Service (NRCS), and USFWS have laws, regulations, policies, and funding programs that authorize and support conservation efforts. In Tooele and Juab counties, some provisions for wildlife or sage-grouse conservation or management are already in place.

### Utah Division of Wildlife Resources (UDWR)

Title 23 of the Utah Code is the Wildlife Resources Code of Utah and provides the UDWR with the powers, duties, rights, and responsibilities to protect, propagate, manage, conserve, and distribute wildlife throughout the state. Section 23-13-3 declares that wildlife existing within the state, not held by private ownership and legally acquired, is property of the state. Sections 23-14-18 and 23-14-19 authorize the Utah Wildlife Board to prescribe rules and regulations for the taking and/or possession of protected wildlife.

The UDWR's wildlife management philosophy is captured in its Mission Statement, Strategic Plan, and Comprehensive Wildlife Conservation Strategy (CWCS) approved in 2005 (also known as the Utah Wildlife Action Plan). The mission of the Division of Wildlife Resources is "...to serve the people of Utah as trustee and guardian of the state's wildlife, and to ensure its future and values through management, protection, conservation and education." There are three goals associated with this mission. The resource goal states that the UDWR intends to, "Expand wildlife populations and conserve sensitive species by protecting and improving wildlife habitat." The UDWR 2005-2015 Strategic Plan calls for focusing efforts on increasing the abundance, distribution, and range for species of conservation need by sustaining and restoring habitat functions. A ten-year, 2005-2015 Comprehensive Wildlife Strategy (a.k.a. Utah Wildlife Action Plan) was approved in 2005 to address species and habitat of greatest conservation need, priorities for conservation, and actions and future implementation opportunities through partnerships.

Sage-grouse are classified as a "State Species of Concern" and are among the terrestrial species identified as being in the second tier (i.e., Tier II) of three priority categories of species identified in the CWCS. Approximately 60 species across 5 taxa in Utah are identified as being potentially petitioned for placement on the ESA defined Threatened and/or Endangered Species list.

## Counties

The Tooele and Juab County Commissions serve as the executive and legislative branches of local government. They have the authority to:

1. Protect and promote the health, welfare, and safety of the people of Tooele and Juab Counties.
2. Regulate land use, land planning, and quality and protection of natural resources.
3. Adopt regulations and policies to exercise such authorities, including the review and approval or denial of proposed activities and uses of land and natural resources.

The Tooele County General Plan (Tooele County 2006) call for the maintenance of open space and preservation of critical wildlife habitat. Specific goals related to protection of wildlife and habitat include:

- To protect native wildlife, development which interferes with wildlife and their habitats should be avoided. Knowledge of wildlife and their habitats will aid in determining designations for appropriate locations and densities of development in those areas.
- The preservation of open space is important to maintain important pristine mountain views, watershed systems, as well as important valley views and general rural character of the County. Open space includes agricultural lands as well as undeveloped hillsides and fields. Land-use plans should result in decreased development pressure on threatened open space and agricultural areas.

The Juab County Zoning Ordinances designate a Grazing, Mining, Recreation, and Forestry District the objectives of which are to:

1. Preserve, insofar as possible, natural scenic attractions, natural vegetation, and other natural features located within the district.
2. Promote tourism, grazing, mining, and the development of natural resources.
3. Promote sanitation and protect and conserve the water supply and other natural resources.
4. Prohibit substandard, urban type developments.
5. Coordinate with programs of public land agencies.

Some forms of development are permitted in this zone (Juab County Planning Commission).

## Natural Resources Conservation Service (NRCS)

The United States Department of Agriculture (USDA) NRCS has authority to conserve sage-grouse through:

1. The Soil Conservation and Domestic Allotment Act of 1936, as amended (P.L. 74-46)
2. The Department of Agriculture reorganization Act of 1994 (P.L. 409-354; 7 U.S.C. 6962)
3. The Farm Security and Rural Investment Act (Farm Bill) of 2002 (P.L. 107-171)

The NRCS and Farm Service Agency (FSA) jointly implement programs which provide landowners with technical and financial assistance to restore and protect grassland, rangeland, pastureland, shrub land, and certain other lands, through long-term agreements and easements.

The USDA NRCS offers help to private landowners through the 2002 Farm Bill programs to

improve their range and pastureland for sage-grouse habitat. These practices include watershed practices on their private lands, such as water developments and fencing for prescribed grazing to improve livestock distribution. Help is also provided for vegetation or brush management practices, with the seeding of introduced and native species of grasses and forbs for forage improvement to benefit both wildlife and domestic animals. Other Farm Bill programs include wildlife enhancement, conservation easements, watershed and riparian programs, and programs to reduce soil erosion.

### Bureau of Land Management (BLM)

The United States Department of Interior (USDI) BLM has authority for conservation of sage-grouse through:

1. The Federal Land Management Policy Act (FLMPA) of 1976 (43 U.S.C. 1701 et seq., 90 stat. 2743; PL 94-579)
2. The Sikes Act, Title II (16 U.S.C. 670 et seq.), as amended
3. The BLM Manual 6840, Special Status Species Management

Specifically, the FLMPA guidance on sensitive species authorizes that “the public lands be managed in a manner that will protect the quality of scientific, scenic, historical, ecological, and environmental, air, and atmospheric, water resource, and archeological values; that, where appropriate, will preserve and protect certain public lands in their natural condition; that will provide food and habitat for fish and wildlife and domestic animals...(43 U.S.C. 1701 Sec. 102 (a) (8)).”

The 6840 Manual defines Special Status Species as “...any species which is listed, or proposed for listing, as threatened or endangered by the U.S. Fish and Wildlife Service or National Marine Fisheries Service under the provisions of the Endangered Species Act; any species designated by the U.S. Fish and Wildlife Service as a ‘listed’, ‘candidate’, ‘sensitive’ or ‘species of concern’, and any species which is listed by the State in a category implying potential danger of extinction.” The Manual provides for the BLM to implement management plans that conserve these species and their habitats, and to ensure that actions authorized, funded, or carried out by the BLM do not contribute to the need for the species to become listed under provisions of the ESA. In addition, the USFWS Policy: State-Federal Relationships (43CFR part 24.4 (c)), contends that the Secretary of the Interior is responsible for the management of non-wilderness BLM lands for multiple uses, including the conservation of fish and wildlife populations. Finally, the BLM provides conservation guidelines for management of sage-grouse on BLM lands in the National Sage-grouse Habitat Conservation Strategy (BLM 2005a).

### School and Institutional Trust Lands Administration (SITLA)

SITLA was created in 1994 to manage twelve real estate trusts granted to Utah at statehood (1896) by the United States federal government. SITLA is an independent agency of the state government established to manage those lands for the support of common schools and other beneficiary institutions, under the Utah Enabling Act (Title 53C-School and Institutional Trust Lands Management Act).

Title to these trust lands is vested in the state as trustee to be administered for the financial support of the trust beneficiaries. As trustee, SITLA must manage the lands and revenues

generated from the lands in the most prudent and profitable manner possible, and not for any purpose inconsistent with the best interest of the trust beneficiaries. The trust principles impose fiduciary duties upon the state, including a duty of undivided loyalty to, and a strict requirement to administer the trust corpus for the exclusive benefit of, the trust beneficiaries. The beneficiaries do not include other governmental institutions or agencies, the public at large, or the general welfare of the state. SITLA must be concerned with both incomes for the current beneficiaries, and the preservation of the trust corpus for future beneficiaries, which requires a balancing of short and long-term interests so that long-term benefits are not lost in an effort to maximize short-term gains. SITLA has no jurisdiction over wildlife populations on trust lands. Management of rangelands is addressed in Section 53C-5-101 of the School and Institutional Trust Lands Management Act, which states: 1) The director is responsible for the efficient management of all range resources on lands under the director's administration, consistent with his fiduciary duties of financial support to the beneficiaries; and 2) This Management shall be based on sound resource management principles.

#### United States Forest Service (USFS)

The USFS has authority for conservation of sage-grouse through:

1. The Multiple-Use Sustained Yield Act (MUSY) of 1960 (P.L. 86-517, 74 Stat. 215, 16 U.S.C. 528, 528-531)
2. The Sikes Act of 1960 (P.L. 86-797, 74 Stat. 1052, 16 U.S.C. 670 et seq., as amended)
3. The Forest and Rangeland Renewable Resources Planning Act (RPA) of 1974 (P.L. 93-378, 88 Stat. 476, as amended; 16 U.S.C. 1600, 1600-1614)
4. The National Forest Management Act (NFMA) of 1976 (P.L. 94-588, 90 Stat. 2949, 16 U.S.C. 472 et seq.) and its implementing regulations (36 CFR 219, 2005)
5. Public rangelands Improvement Act of 1978 (P.L. 95-514, 92 Stat. 1806, 43 U.S.C. 1901-1908)
6. USDA Regulation 9500-4 and the Forest Service Manual (FSM) Chapter 2600

MUSY directs the USFS to administer the National Forest for multiple uses including fish and wildlife purposes, in cooperation with interested state and local governmental agencies, and others. 'Multiple use' refers to the congruent and coordinated management of the various surface renewable resources so that they are utilized in a manner that will best meet the needs of the American people. The Sikes Act provides authority for cooperative planning, habitat improvement, and providing adequate protection for species considered to be threatened, rare, or endangered by a State agency. RPA and NFMA provide for comprehensive, integrated planning that will provide for the diversity of plant and animal communities to meet overall multiple-use objectives. USDA Regulation 9500-4 directs the USFS to manage "habitats for all existing native and desired nonnative plants, fish, and wildlife species in order to maintain at least viable populations of such species." USFS policy includes provisions for the development of conservation strategies for species that could be negatively affected by forest plans or proposed projects (FSM 2621.2).

#### Memorandum of Understanding

There are two Memoranda of Understanding (MOU) that address conservation of sage-grouse. The first was signed in 1999 by members of the Western Association of Fish and Wildlife Agencies (WAFWA) to promote conservation and management of sage-grouse and their

habitats. Thirteen states, including Utah, and two Canadian provinces were signatories to that MOU. The second MOU, signed in 2000, is between WAFWA, USFS, BLM, and the USFWS. This MOU provides for cooperation among state, provincial, and federal agencies in the development of a rangewide strategy to direct conservation of sage-grouse and their sagebrush habitats.

A MOU between state and federal agencies within the state of Utah is currently being developed. The MOU promotes the conservation of sage-grouse and their sagebrush habitats, encourages cooperation between signatories, and supports Adaptive Resource Management Local Working Groups as the primary format for addressing sage-grouse and sagebrush steppe issues in the state.

### **G. Policy for Evaluation of Conservation (PECE) Standards**

The PECE Standards set criteria for the USFWS to use in determining whether a formalized conservation effort contributes to making listing a species unnecessary or contributes to forming a basis for listing a species as threatened, rather than endangered. The draft PECE was published on June 13, 2000 (65 FR 37102), and was finalized on March 28, 2003 (68 FR 15100-115). The PECE contains nine criteria the USFWS will use to evaluate that the conservation effort will be implemented, and six criteria to determine if the action will be effective. Conservation efforts included under this policy include those identified in conservation agreements, conservation plans, management plans, or similar documents developed by federal agencies, state and local governments, tribal governments, businesses, organizations, individuals, and a combination of the above. The criteria are not considered comprehensive. The USFWS will consider all appropriate factors and unique, specific circumstances when evaluating formalized conservation actions.

PECE reviews will be conducted to individual conservation actions (rather than conservation plans). Should Greater Sage-grouse be petitioned for listing or be listed under the ESA, this Plan will be reviewed and assessed as part of the preparation of a listing decision and will follow the most recent procedural guidance.

### III. Conservation Assessment

#### A. General Sage-grouse Biology/Ecology

Numerous authors have described various aspects of sage-grouse biology, ecology, and life history. In recent years, several more have published summaries. For the purposes of this document, we have included the summary from the Statewide Strategic Plan (UDWR 2002) and would also recommend the Conservation of Sage-grouse and Sagebrush Habitats by Connelly et al. (2004) for a thorough discussion.

##### Physical Description

The sage-grouse is the largest grouse species in North America. Adult males are larger than adult females. Adult males weigh 4–7 pounds (1.7–2.9 kg) and are 27–32 inches (65–75 cm) long. In comparison, adult females weigh 2–4 pounds (1.0–1.8 kg) and measure 20–25 inches (50–60 cm) long. Both sexes have narrow, pointed tails and a variegated pattern of grayish brown, buff, and black on the upper parts of the body and a diffuse black abdominal pattern. Males have blackish brown throats, a dark V-shaped pattern on the neck, and white breast feathers. When strutting, males inflate two gular sacs of olive green skin and erect hair-like black feathers (filoplumes) on the back of the neck. Females lack the V-shaped pattern, their throats are buff, and their lower throats and breasts are barred with blackish brown (Schroeder et al. 1999).

There are noticeable morphological differences between Greater Sage-grouse and Gunnison Sage-grouse. Gunnison Sage-grouse are two-thirds the size of Greater Sage-grouse. Gunnison Sage-grouse tail feathers have horizontal white barring along their length compared to the variegated pattern found in Greater Sage-grouse. The filoplumes, found only on male sage-grouse, are much thicker and denser in Gunnison Sage-grouse than in Greater Sage-grouse. There are also noticeable differences in the strutting behavior of the two sage-grouse species (Young et al. 2000).

##### Seasonal Movements and Home Range

Sage-grouse populations can be defined as one of two types: 1) non-migratory – grouse do not make long-distance movements between seasonal ranges, and 2) migratory – grouse make long-distance movements between distinct seasonal ranges. Movements between seasonal ranges can exceed 45 miles (75 km; Connelly et al. 1993).

Home range size for migratory sage-grouse populations can exceed 540 mi<sup>2</sup> (1,500 km<sup>2</sup>; Hulet 1983). For non-migratory sage-grouse populations, home range size varies from 4–11 mi<sup>2</sup> (11–31 km<sup>2</sup>). Sage-grouse exhibit high fidelity to seasonal ranges (Fischer et al. 1993). Females return to the same area to nest each year and may choose nests near their previous year's nesting site (Bunnell et al. 2000, Gates 1983).

##### Breeding

The center of breeding activity for sage-grouse is the 'lek' or strutting ground. Male sage-grouse

begin to congregate on leks in early March and perform a ritualized courtship display. Use of leks may continue as late as early June. Mating occurs on the lek. Fifty to ninety percent of the males use leks during the breeding season. As sage-grouse populations decline, the number of males attending leks may decline, or the use of some leks may be discontinued. Likewise, as populations increase, male attendance on leks may increase and/or new leks may be established or old leks reoccupied (Connelly et al. 1981).

### Nesting/Reproduction

Nesting generally takes place one to two weeks after mating and may continue as late as early June (Wallestad 1975). Sage-grouse generally have lower reproductive rates and higher survival rates than other species of upland game birds (Connelly and Braun 1997). Nesting rates vary from year to year and from area to area (Bergerud 1988, Connelly et al. 1993, Schroeder 1997, Coggins 1998). Connelly et al. (1993) reported that in Idaho up to 45% of yearling and 22% of adult female sage-grouse do not nest each year. Schroeder (1997) found that essentially all female sage-grouse in Washington nested. The variation is most likely a result of the quality of nutrition available and the health of pre-laying females (Barnett and Crawford 1994). Re-nesting by sage-grouse varies regionally from 20% (Hulet 1983, Connelly et al. 1993) to greater than 80% (Schroeder 1997). In summary, sage-grouse have the lowest reproduction rate of any North American game bird. As a result, populations are not able to recover from low numbers as quickly as those of most other game birds.

Sage-grouse nest success varies from 12–86% (Trueblood 1954, Gregg 1991, Schroeder et al. 1999). Adult females may experience higher nest success rates than yearling females (Wallestad and Pyrah 1974). However, differential nest success between age groups has not been observed in other studies (Connelly et al. 1993, Schroeder 1997). Nest success is dependent on vegetation cover type (Gregg 1991). Gregg (1991) reported that the highest nest success occurred in mountain big sagebrush (*A. t. vaseyana*) cover type. The greater cover of medium-height shrubs with grass 7 inches (>18 cm) in height, increases sage-grouse nest success (Gregg et al. 1994).

Clutch size of sage-grouse is extremely variable and relatively low compared to other species of game birds (Schroeder 1997). Average clutch size for first nests varies from 6.0–9.5 throughout the species range (Schroeder 1997, Sveum et al. 1998). These differences may be related to habitat quality and overall health of pre-laying females (Coggins 1998).

### Survival Rates

Annual survival rates for yearling and adult female sage-grouse vary from 35–85%; adult male survival rates vary from 38–54% (Wallestad 1975, Zablan 1993, Connelly et al. 1994). Lower survival rates for males may be related to physiological demands of sexual dimorphism and higher predation rates on males during the breeding season (Swenson et al. 1987).

Sage-grouse predators include raptors, coyotes, ravens, squirrels, and skunks. The increase in urban development has resulted in the addition of nonnative predators such as dogs, cats, and foxes (Connelly et al. 1991).

Little information has been published on mortality of juvenile sage-grouse, or the level of

production necessary to maintain a stable population. Among western states, long-term juvenile to hen ratios have varied from 1.40–2.96 juveniles per hen in the fall. In recent years, this ratio has declined to 1.21–2.19 juveniles per hen (Connelly and Braun 1997). It is thought that at least 2.25 juveniles per hen should be present in the fall population to allow for stable to increasing sage-grouse populations (Connelly and Braun 1997, Edelman et al. 1998).

## **B. Habitat Requirements**

### Breeding/Nesting Habitat

Leks, or strutting grounds, tend to be traditional. In general, the same areas are used year after year. Leks typically occur in open areas surrounded by sagebrush (Patterson 1952, Gill 1965). Examples of lek sites include landing strips, old lakebeds or playas, low sagebrush flats, openings on ridges, roads, cropland, and burned areas (Connelly et al. 1981, Gates 1985). Sage-grouse males appear to form leks opportunistically at sites within or adjacent to potential nesting habitat. The lek is considered to be the center of year-round activity for non-migratory grouse populations (Eng and Schladweiler 1972, Wallestad and Pyrah 1974, Wallestad and Schladweiler 1974), but this may not be the case for migratory populations (Connelly et al. 1988, Wakkinen et al. 1992). Average distances between nests and the nearest leks vary from 0.6–3.9 miles (1.1–6.2 km), however, some females may nest more than 12.5 miles (20 km) away from the lek (Autenrieth 1981, Wakkinen et al. 1992, Fischer 1994, Hanf et al. 1994).

Habitat used by pre-laying hens is also part of the general breeding habitat. These areas provide hens with forbs that are high in calcium, phosphorus, and protein, all of which are necessary for egg production. The condition and availability of these areas is thought to have a significant effect on reproductive success (Barnett and Crawford 1994, Coggins 1998).

Most sage-grouse nests are located under sagebrush plants (Patterson 1952, Gill 1965, Gray 1967, Wallestad and Pyrah 1974), however, nests have been found under other plant species (Griner 1939, Connelly et al. 1991, Gregg 1991). Sage-grouse that nest under sagebrush experience a higher nest success than those nesting under other plant species (Connelly et al. 1991). Research on sage-grouse nesting habitat has documented that sage-grouse tend to select sites under sagebrush plants that have large canopies. The canopies provide overhead cover and an herbaceous understory, thus providing lateral cover and allowing birds to be hidden from view (Patterson 1952, Gray 1967, Klebenow 1969, Wallestad and Pyrah 1974, Wakkinen 1990, Gregg 1991, Fischer 1994, DeLong et al. 1995, Bunnell et al. 2000). Herbaceous cover associated with nest sites may provide scent, visual, and physical barriers to potential predators (DeLong et al. 1995).

### Brood-rearing Habitat

Early brood-rearing habitat generally occurs relatively close to nest sites, but movements of individual broods may be highly variable (Connelly 1982, Gates 1983). Early brood-rearing habitats may be relatively open (14% canopy cover) stands of sagebrush when compared to optimum nesting habitat (Martin 1970, Wallestad 1971), but need greater than 15% canopy cover of forbs and grasses (Sveum et al. 1998, Bunnell et al. 2000). High plant species richness with abundant forbs and insects characterize brood areas (Dunn and Braun 1986, Klott and Lindzey 1989, Drut et al. 1994a, Apa 1998). Insects, especially ants and beetles, are an important food component of early brood-rearing habitat (Drut et al. 1994a, Fischer 1996). As herbaceous plants mature and dry, hens usually move their broods to more mesic sites during June and July where more succulent vegetation is available (Gill 1965, Klebenow 1969, Connelly et al. 1981, Connelly et al. 1988, Fischer 1996, Bunnell et al. 2000). Sage-grouse broods occupy a variety of habitats during summer including sagebrush, relatively small burned areas within sagebrush, wet meadows, farmland, and other irrigated areas adjacent to sagebrush habitats (Savage 1969,

Martin 1970, Connelly et al. 1981, Gates 1983, Connelly et al. 1988, Pyle and Crawford 1996).

Late brood-rearing habitats are highly variable. Patterson (1952) reported that grouse move from summer to winter range in October but during mild weather in late fall, some birds may still use summer range. Movements from fall to winter ranges are slow and meandering, and occur from late August to December (Connelly et al. 1988). Wallestad (1975) documented a shift in feeding habits from September, when grouse were consuming a large amount of forbs, to December when birds were feeding only on sagebrush.

#### Winter Habitat

Sage-grouse winter habitats are relatively similar throughout most of their range. Because their winter diet consists almost exclusively of sagebrush, winter habitats must provide adequate sagebrush that is accessible through the winter. Eng and Schladweiler (1972) and Wallestad (1975) indicated that most observations of sage-grouse during winter in Montana occurred in sagebrush habitats with greater than 20% canopy cover. However, Robertson (1991) indicated that sage-grouse used sagebrush habitats that had average canopy cover of 15%. Sage-grouse tend to select areas with both high canopy cover and taller big sagebrush (*Artemisia tridentata*).

During winter, sage-grouse feed almost exclusively on leaves of sagebrush (Patterson 1952, Wallestad 1975). Big sagebrush dominates the diet of sage-grouse in most portions of their range (Patterson 1952, Wallestad 1975, Remington and Braun 1985, Welch et al. 1988) but low sagebrush (*A. arbuscula*), black sagebrush (*A. nova*; Dalke et al. 1963, Beck 1977), fringed sagebrush (*A. frigida*; Wallestad 1975), and silver sagebrush (*A. cana*; Aldridge 1998) are also consumed in many areas depending on the availability. Sage-grouse in some areas apparently prefer Wyoming big sagebrush (*A. t. wyomingensis*; Remington and Braun 1985, Meyers 1992) and in other areas mountain big sagebrush (*A. t. vaseyana*; Welch et al. 1988). Some of the differences in selection may be due to preferences for higher levels of protein (Remington and Braun 1985).

It is critical that sagebrush be exposed at least 10–12 inches (25 cm) above snow level (Hupp and Braun 1989). This provides both food and cover for wintering sage-grouse. In situations where snow covers the sagebrush, birds will move to areas where sagebrush is exposed.

During winter, sage-grouse will either partially or completely bury themselves in snow (snow roosting) for added thermal protection from winter temperatures.

### **C. Distribution and Abundance**

Populations of Greater Sage-grouse have been declining for the past 25 years (Braun 1995, Connelly and Braun 1997, Beck et al. 2003, Connelly et al. 2004). Concerns about population status and distribution have heightened awareness about the appropriateness of various monitoring efforts and techniques. Connelly et al. (2000) indicated that monitoring was a key component of sage-grouse management. Utah's Strategic Management Plan (UDWR 2002) also emphasizes the need to monitor sage-grouse populations and habitats. Further, the MOU signed by WAFWA representatives in 1999 with federal agencies (2000) calls for consistent monitoring and data collection.

Several techniques have historically been used in Utah and in the Resource Area to assess sage-grouse population trends, status, and distribution including lek counts, brood surveys, field bag checks, wing barrels, and hunter surveys. Currently, the primary technique employed by biologists in Utah and in the Resource Area is lek counts. This method is described in detail later in this section.

#### Historic Distribution of Sage-grouse

Determining historic distribution of sage-grouse is difficult and problematic for several reasons, but primarily because scientific studies are not available from the historic time frames in question. For many areas, no written or zoological records exist. It is thought that sage-grouse once existed in all 29 Utah counties. Today sage-grouse are found in 26 counties in Utah and are thought to occupy 50% of the habitat they once did (UDWR 2002).

The Rangewide Conservation Assessment of Greater Sage-grouse and Sagebrush Habitats (hereafter referred to as the Rangewide Plan; Connelly et al. 2004) analyzed the historic distribution of sage-grouse based on sage-grouse distribution maps, habitat maps, sage-grouse museum specimens, known lek locations, and research on sage-grouse movement patterns and habitat use. The authors define historic distribution as a 'pre-settlement' distribution, occurring prior to 1800 when rapid settlement by people of European descent began.

The Rangewide Plan describes potential historic distribution of sage-grouse in the West Desert as part of an evaluation of historic range in the Southern Great Basin (Connelly et al. 2004). The authors suggest that there are three apparently separate areas within the Southern Great Basin. The Resource Area is associated with what is described as "...a relatively continuous area in central and eastern Nevada and the western edge of Utah," and, "...a fragmented area associated with the north-south oriented mountain ranges in central Utah." The authors further suggest that although barren habitats west of the Great Salt Lake and forested and alpine areas in mountainous areas were not historically occupied by sage-grouse, the distribution of sage-grouse in this area is still reduced from pre-settlement times (Connelly et al. 2004).

#### Lek Counts

During the breeding season, sage-grouse congregate on a relatively small number of sites, called leks, to display and breed. Because sage-grouse demonstrate high fidelity to lek sites, they offer the best opportunity for monitoring populations (Jenni and Hartzler 1978, Beck and Braun 1980,

Connelly et al. 2000, 2003, 2004). Lek count methodology was first described by Patterson (1952), who studied sage-grouse in Wyoming, and was based on a count of the maximum number of males observed on a lek over a series of 3–4 visits. The method described by Patterson (1952) results in an index of the population. Population indices, commonly used by wildlife managers, involve a count, or measurement, of some aspect of the population that is both convenient to measure and thought to be related to abundance (e.g. bird calls, pellet counts, roadside observations, track surveys). Anderson (2001), whose primary criticism was that they fail to lead to defensible estimates of population size or status, described the shortcomings of this type of sampling. With regards specifically to lek counts, Beck and Braun (1980) noted that they only lead to conclusions about population size and status when the following information is known: total number of leks in an area, attendance patterns of adult and yearling males, inter-lek movements patterns, and the ratio of males to females (i.e. the relationship between the maximum count and the size of the population).

Despite the problems associated with indices and lek counts specifically, they remain the best and primary available means for assessing population trends and estimating population size and status (Autenrieth et al. 1982, Connelly et al. 2000). Throughout Utah, lek counts are conducted between late February and May (depending on weather conditions and access to lek sites) on all known leks to the greatest extent possible. Leks are counted three to four times during this period and counts are made between 30 minutes before and 1 hour after sunrise. An estimate of population size is calculated based on the following assumptions: 1) 75% of all males were counted on strutting grounds, and 2) that the male:female ratio in the population is 1:2 (UDWR 2002).

The number of active leks in an area can also be used as an indicator of population size. Cannon and Knoph (1981) noted that lek numbers seem to increase roughly in proportion to population size. There is evidence that as population size increases, established, ‘traditional’ lek attendance increases and smaller ‘satellite’ leks appear and then disappear as population size decreases again. In Utah, a lek is defined as a site or area traditionally used for display. Leks are considered ‘active’ when at least two males have been observed for at least three years. Conversely, leks are considered ‘inactive’ when birds have been absent from a traditional site for more than three years. The use of the number of ‘active’ or ‘traditional’ leks as an indicator of population size is also problematic. Satellite leks are typically smaller and are likely to be less noticeable, lek detection is likely to vary with both density of leks and population density, and search effort likely plays a large role in detection and consistency of measurement.

**D. Assessment of Local Population**

Plan Area

The West Desert Resource Area is located in Tooele and Juab counties in western Utah (Figure 1). The Resource Area encompasses 5,137,991 acres and is divided into two subunits, Vernon and Ibapah, according to sage-grouse population distribution. The Resource Area is bounded on the south by the Juab County-Millard County line, on the east by Tooele County-Utah County boundary and Highway 6, on the north by I-80, and on the west by the Utah-Nevada border, excluding land managed by the U.S. Department of Defense. The Resource Area is managed primarily by the USFS, BLM, and private landowners. The predominant land use in the area is grazing by domestic livestock.

The West Desert is characterized by hot summers and cold winters. According to National Climate Data Center records, temperatures range from an average high of around 90° F in July to an average low of about 12° F in January. As the name implies, the West Desert is a dry region of the state. Ibapah receives an average of only 9.74 inches of annual precipitation; Vernon receives slightly more with an average of 10.52 inches. Most precipitation comes in the form of snow during January.

*Landownership*

Most of the Resource Area is public land with smaller areas managed by the state of Utah, the USFS, and private landowners (Figure 1, Table 2).

Table 2. Landownership in the WDARM Resource Area.

Subunit	Landowner	Area (acres)*
Deep Creek	BLM	501,683
Deep Creek	Department of Defense	2,013
Deep Creek	Tribal	93,183
Deep Creek	Private	35,461
Deep Creek	State Trust	34,669
Vernon	BLM	498,233
Vernon	Department of Defense	43,985
Vernon	Tribal	9,558
Vernon	Private	386,159
Vernon	State of Utah	956
Vernon	US Forest Service	179,085
Vernon	State Trust	92,949
Great Salt Lake Desert	BLM	6,941,504,024
Great Salt Lake Desert	USFWS	14,917
Great Salt Lake Desert	Tribal	8,582
Great Salt Lake Desert	State Trust	203,763
Great Salt Lake Desert	State of Utah	5,096
Great Salt Lake Desert	Private	183,598
Great Salt Lake Desert	Department of Defense	1,367,688
*Water accounts for 67,825 acres (1.24%) of the total acreage of the Resource Area.		




Figure 1. The WDARM Resource Area with land management and landownership designated.

### *Wildlife Populations*

Several species of birds, small mammals, and reptiles are found only in sagebrush environments. Passerine birds obligated to use sagebrush environments include Brewer's sparrow, sage sparrow, and sage thrasher. Additionally, though not obligated to use only sagebrush environments, vesper sparrow and loggerhead shrike are commonly found in sagebrush communities in the Resource Area. Other obligate species include the sagebrush vole and the sagebrush lizard. In addition to these obligates, a large number of other birds, small mammals, and reptiles commonly make use of sagebrush environments within the Resource Area.

While sage-grouse populations in the Resource Area have been counted and studied, little or nothing is known about the local status of these other wildlife species listed above. It is assumed that their numbers and geographic extent are tied to the condition and extent of big sagebrush communities. This plan operates with the intent that maintenance of substantial areas of high quality sagebrush steppe, measured by healthy populations of sage-grouse, will provide sufficient habitat for these other sagebrush obligate species to thrive in the Resource Area.

In parts of the Resource Area, sage-grouse populations overlap with populations of pygmy rabbits (*Brachylagus idahoensis*). The distribution, behavior, and ecology of this species is currently under investigation by researchers from Brigham Young University (BYU; Larsen et al. 2006). Information obtained from this research will be used to help refine ecosystem and community-level conservation and management to help maintain populations and habitats for both pygmy rabbits and sage-grouse.

### *Human Populations*

Prehistoric Indian sites from the Desert Archaic and Fremont Indian cultures have been discovered in several areas of the West Desert indicating that human habitation of the area has occurred for over 10,000 years. The Goshute Indian Tribe claims the West Desert as their ancestral home. Today, the tribe has a reservation in Skull Valley and in the Deep Creek Watershed, with tribal headquarters located in Ibapah (Marsh 2005).

Early explorers to the area included the Dominguez-Escalante expedition, Jedediah Smith, John W. Gunnison, J. H. Simpson, and John C. Fremont. Explorations by Gunnison and Simpson later lead to the establishment of the Pony Express route and transcontinental telegraph (Utah State Historical Society 1998).

The first permanent immigrant settlements were established in the mid 1800s by Mormon Pioneers. Three short-lived military outposts were established in the area; one at present day Stockton and one at Fairview in Cedar Valley. Mining during the 1860s was responsible for accelerating early growth. Several towns were established in the Oquirrh Mountains where silver, lead, zinc, and gold were extracted. The mining industry attracted a diverse population to Tooele Valley and also brought the railroad to the area, further increasing opportunities for growth (Marsh 2005, Utah State Historical Society 1998).

Military installations established during World War II also boosted Tooele County's population and brought millions of dollars to the local economy. Today the Tooele Army Depot and

Dugway Proving Grounds occupy much of the West Desert and employ many county residents (Marsh 2005).

In Tooele County, human populations are located in Tooele Valley (Tooele, Grantsville, Stansbury Park, Erda, Pine Canyon, and Lakepoint), Rush Valley (Stockton, St. Johns, Clover, Ophir, and Vernon), Skull Valley, (Dugway, Terra, and Iosepa), and on the Utah/Nevada border at Wendover and Ibapah. Although growth is projected for all parts of Tooele County, it is likely to target areas closest to the Wasatch Front (Cline 2006); some of these areas include sage-grouse habitat. Juab County has grown approximately 40% in the past decade (U.S. Census Bureau as interpreted by Negative Population Growth 2006). This growth is projected to continue and be concentrated in the eastern part of the county, around the city of Nephi.

### *Livestock Grazing*

Except for a handful of passing explorers and immigrants, the Tooele Valley was first used as pasture by members of the Church of Jesus Christ of Latter Day Saints. Three days after the first company of Mormon pioneers arrived in the Salt Lake Valley (July 24, 1847), an exploring party ventured west to the southern shores of the Great Salt Lake and viewed Tooele Valley from the northeast corner. The large valley with abundant grass was obviously well suited for grazing which began the same year. In 1849, Howard Stansbury established a post at Adobe Rock for the purpose of surveying the Great Salt Lake. The government mules and livestock were tended by herders who built the first house in the valley. As the number of livestock increased in the territory, more and more pasture was needed. In the fall of 1849, over 300 head were brought to the valley to winter (Blanthorn 1998). Permanent settlements were soon established in Tooele, Rush, and Skull Valleys; all centered on grazing. Because of the excellent pastures in the county, especially for winter grazing, livestock numbers increased until the range began to deteriorate. By the end of the 1870s, conditions were such that some of the best range in the region had become some of the worst (Blanthorn, 1998). Numbers began to decrease toward the end of the century, but large bands of sheep trailing through the county continued to deplete the range. It is reported that approximately one-half million sheep trailed through the Tooele Valley at the turn of the century (Blanthorn, 1998). In 1934, conditions were right to create what became known as the Grantsville Dust Bowl. A combination of factors including restricted grazing, reseeding, the Taylor Grazing Act, the establishment of the Tooele Army Depot and the Grantsville Soil Conservation District brought about range improvement. Since then there have been episodes of drought and infestations of Mormon Crickets, but nothing as serious as the dust bowl. The number of livestock grazing in the county has remained relatively constant since initial grazing improvements were made. As of 2006, the National Agricultural Statistics Service reported 15,000 head of beef cattle and 27,000 head of total cattle in the county (USDA-NASS, 2006).

### *Farming*

Agricultural production in the Resource Area began with early settlers. Agricultural farms were, and still are, present throughout Tooele County where primarily hay and other field crops are produced. The first lucern seed was brought to Tooele County in 1863 by James James.

### Population Status and Distribution

Sage-grouse are believed to have existed as a species in North America for approximately 350,000 years. Sage-grouse likely were historically found in all 29 Utah counties and abundant where suitable habitat existed throughout Utah until the early 1900s (Beck et al. 2003). Today, sage-grouse are found in 26 of Utah's counties and are thought to occupy only 50% of the geographic area they once did (Beck et al. 2003).

The UDWR began monitoring sage-grouse populations in the Resource Area by annually counting males on leks in 1968 and 1982, in the Vernon and Deep Creek Subunits, respectively (Figure 2). Subunits are evaluated separately because there is likely no movement between the two areas (Robinson, unpublished data). When monitoring began in the Vernon Subunit, a total of 44 male sage-grouse were counted on two leks. In 1982, 20 males were counted on one lek in the Deep Creek Subunit. The Vernon Subunit high count was recorded in 2002 when 163 males were counted on six leks. Under the assumption that 75% of all males in the population were observed and counted, and assuming a sex ratio of 1.67 females to each male (Zablan 1993), the estimated spring population size in the Vernon Subunit was approximately 326 adult birds in 2002. New leks discovered in recent years will likely result in a new high count in the Vernon Subunit as monitoring continues and these new leks are considered active and included in indices. New leks are not considered active until at least two males are observed for two years. Further, population estimates based on lek counts should be treated cautiously due to variance in the methods used to collect lek count data, the assumptions built into the estimate, and other factors. However, as no other population estimation technique is currently available, WDARM will use this currently established method. There is no high count available for the Deep Creek subunit because leks have not been monitored consistently in this area.

In 2005, a total of 143 males were counted on two known active and one new lek in Vernon. In Ibapah, a total of 59 males were counted on one known active, and two new leks. In 2006, two additional new leks were discovered and a total of 190 males were counted on six total leks. Also in 2006, a total of 93 males were counted on five total leks, one of which was discovered that year.

An observation of the number of males per lek is another index used to evaluate sage-grouse population trends. Because this index accounts for the number of leks counted (i.e. the amount of effort) this index may, in cases where effort is variable, be a more useful illustration of the population trend. In the Vernon Subunit, the number of males per lek still reflects a variable but stable pattern in sage-grouse numbers since the late 1960s (Figure 3).

It appears that population monitoring through the use of lek counts has been somewhat inconsistent in the past, although increased efforts in 2005 and 2006 have resulted in six new leks being discovered. As Figures 2 and 3 illustrate, there are years when no counts were made. Lek sites can be difficult to access in some years due to inclement weather and road conditions. Additionally, leks may be located on private or Tribal land and permission to access them may not be available.




Figure 2. Maximum total number of males counted on all leks in the Vernon Subunit of the Resource Area, 1968-2006.




Figure 3. Number of males per lek for the Vernon Subunit of the Resource Area, 1982-2006.

## **Local Ecology and Life History**

Little published information is available regarding the ecology and life history of sage-grouse populations in the Resource Area. Although monitoring (lek counts) has been conducted somewhat regularly since the late 1960s (Figures 2 and 3), few studies have documented information about aspects of habitat use, survival, sources of mortality, and reproductive success. Since 2005, sage-grouse in both subunits of the Resource Area have been studied by a graduate student under the direction of Dr. Terry Messmer at USU. Research has focused on the local ecology, life history, movement, sources of mortality, and habitat use of sage-grouse in both the Vernon and Deep Creek Subunits. Rather than summarizing research conducted in the Resource Area, we have elected to provide all available quarterly reports, annual reports, and final reports for the study in Appendix B. Research efforts in and around the Resource Area are ongoing and will be used continuously to revise and adapt management strategies and this Plan.

### Local Habitat

In 1999 and again in 2006, the UDWR mapped the extent of seasonal habitat types in the Resource Area. Figures 4 and 5 illustrate where nesting, brood-rearing, and winter habitats are located in the Resource Area.

The UDWR Big Game Range Trend project has been monitoring sites throughout the Resource Area to track changes in vegetation composition, structure, and diversity. Although these sites were placed in areas used by big game, where they overlap with sage-grouse seasonal habitat types (Figure 6), they can still provide some information about vegetation and habitat conditions in those areas in a general sense. Data collected at these sites are summarized and available at: <http://www.wildlife.utah.gov/range/>.

### Habitat Improvements and Completed Conservation Actions

Several habitat improvement projects in the Resource Area have been implemented by WDARM partners and were targeted at restoring or enhancing sage-grouse habitat. Treatments were generally aimed at reducing sagebrush canopy and enhancing native grass/forb cover in the understory. Additional habitat improvement projects were planned for 2006. The UPCD state and regional teams are currently addressing habitat issues with their statewide watershed initiative which focuses on the protection, management, and/or restoration of important sagebrush-steppe habitats. The UPCD is made up of a variety of partners including state and federal land management agencies, private landowners, universities and extension services, soil conservation districts, and county and local entities. The Central Region UPCD team has delineated focus areas within the Resource Area based upon critical sage grouse habitats, and is currently working on identifying projects and acquiring funding to implement restoration activities. Habitat restoration projects involving the reduction of expanding pinyon-juniper forests into sagebrush habitats have already begun in the Vernon subunit. Likewise, a project to enhance sage grouse wintering habitat on BLM lands was completed in the Deep Creek subunit in 2005. Several Big Game Range Trend sites were established in 2006 to monitor treatments. Most of these projects have been a combination of fence, water development, fuels reduction projects, and brush management. The locations of some projects conducted in the Resource Area are illustrated in Figure 7; acreage of past and proposed treatments is listed in Table 3.




Figure 4. Location of sage-grouse nesting and brood-rearing habitat in the WDARM Resource Area, as identified by the UDWR, 2006.




Figure 5. Location of sage-grouse winter habitat in the WDARM Resource Area, as identified by the UDWR, 2006.




Figure 6. Location of Utah Big Game Range Trend Sites and vegetation monitoring sites located within sage-grouse seasonal habitat, as identified by the UDWR, 2006.




Figure 7. Location of some habitat improvement projects implanted by various WDARM partners on public and private land in the Resource Area.

Table 3. Habitat improvement projects implemented in 2004, 2005, 2006, and proposed for 2007 by the WDARM partners on public and private land in the Resource Area.

Year	Project Name	Description	Acres
2005	Middle Pasture	Aerator, broadcast, and aerial seeding	1000
	Clover Creek	Bullhog, aerial seeding	400
	Iosepa	Bullhog, aerial seeding	400
	Bennion Ranch	Lop and scatter	150
2006	Sage Valley	Harrow, broadcast seeding	500
	Goshute Chaining	2-way chaining, aerial seeding	800
	Bennion Ranch	2-way chaining, aerial seeding	500
	Bennion Ranch	Spike	160
	St. John	Aerial seed, 1-way chain	1200
	East Onaqui	Harrow, broadcast seeding	200
2007 (funded)	East Onaqui	Bullhog, aerial seeding	500
	Big Hollow	Bullhog, aerial seeding	600
2007 (proposed)	Sage Valley	Lop and scatter	1300
	Clover Creek	Chaining and harrow, reseeding	550
	Ibapah	Harrow, broadcast seeding	250
	East Pasture	Harrow, broadcast seeding	150
	Stansbury Mtns. West slope	Bullhog, Rx burn	60
	Spanish Fork District	Noxious weed control	650
	Big Hollow	PJ thinning, reseeding	55

#### IV. Threat Analysis

In this section, we summarize and describe the potential threats to sage-grouse populations in the Resource Area. Where possible, we describe actual, known impacts to sage-grouse and their habitats; however, due to a lack of empirical information regarding many of the threats described, we are only able to present general information and make educated extrapolations about the local area. Potential threats are listed in alphabetical order below.

##### A. Altered Water Distribution

Water diverted into pipes from canals and ditches to improve flood control and irrigation efficiency has altered natural water sources. Lining ditches with concrete have made them less permeable. Some streams and rivers have been channelized to improve water distribution for irrigation and agricultural efficiency. These changes have also resulted in a loss of ground water, riparian habitat, and mesic/wet meadow sites. Channelization has resulted in increased bank erosion in some areas, and downcutting effects in both small creeks and larger streams and rivers.

In YEAR, the Vernon Watershed Plan was published. The primary focus of the Vernon Watershed Plan was the installation of the Vernon Reservoir and associated pipelines to private lands and irrigation systems. The Vernon Watershed Plan also states that sage-grouse are an important game species and warrant management. Specifically, it states: “Sage grouse populations have suffered a serious decline over the years. Restoration of this species to huntable numbers is considered the primary game management problem. Lack of suitable habitat appears to be one of the principal limiting factors, and attention should be directed toward improving habitat for sage-grouse wherever possible.”

Wet meadows and mesic sites provide succulent vegetation for sage-grouse and sage-grouse broods, especially in the drier late summer months. These areas are also an important source of insect food resources for both chicks and adult sage-grouse. Mesic sites seem to be more prevalent in the Deep Creek Subunit and may be lacking in the Vernon Subunit. Ongoing research will address issues related to habitat use and availability in the entire Resource Area. Further research may be warranted.

Comment [kw1]:

## **B. Development and Human Infrastructure**

In this section, we summarize the potential effects of development and human infrastructure including: 1) homes and cabins, 2) power lines, fences, and other tall structures, 3) renewable and non-renewable energy, and 4) roads, on sage-grouse populations in the Resource Area. These impacts were considered together because they are associated with similar stresses (loss of habitat quality and quantity, habitat fragmentation, direct disturbance, increased predator pressure). There is little empirical evidence available regarding the direct or indirect impacts of most of the threats reviewed in this section, especially that are specific to the Resource Area.

### *Home and Cabin Development*

Home and cabin development impacts sage-grouse populations through direct loss of habitat, habitat fragmentation, increased domestic predation (i.e. dogs and cats), and can lead to increases in other threats including power lines, fencing, roads, and incompatible OHV recreation.

Increases in the human population in and around the Resource Area have led to corresponding increases land development. Cities in the Resource Area are increasingly becoming bedroom communities for Salt Lake City.

Increases in the human population in the West Desert have led to corresponding increases in the amount of land being developed for homes and industry. Residential development in the Resource Area reached a peak in the early 1990s and in recent years has declined somewhat with about 450 and 45 new building permits issued each year in Tooele and Juab Counties, respectively (Bureau of Business and Economic Research, University of Utah 2006). Most housing development in the West Desert is associated with existing towns and cities, especially the towns of Tooele, Stansbury, and Eureka and likely does not directly impact areas where sage-grouse are known to occur.

### *Power lines, Fences, and Other Tall Structures*

Although this threat is poorly understood, sage-grouse are potentially subject to increased mortality and disturbance resulting from man-made structures including fences, power lines, and other tall structures (wind turbines, communication towers). Sage-grouse may fly into these structures which can result in death or may injure them to the point where they can not effectively avoid predators. Sage-grouse mortalities due to collision with power lines, fences, and other tall structures have been observed in Colorado, Utah, and other areas (Gunnison Sage-grouse Rangewide Steering Committee 2005). Construction of any structure can result in some habitat loss and fragmentation. Fragmentation may increase vulnerability to predation. Fences have increased in number over the years, as allotments have been split and cross-fenced with the development of rural properties, and with construction of new county roads. Power lines have also increased in number and length (Figure 8), and transmission and service lines have been constructed to service mines and transfer electric power out of the area.

*Roads*

Collisions with motor vehicles, either while flying or while walking on or across roadways, are other potential causes of direct mortality or severe injury for sage-grouse. Road construction can cause an increase in dust on plants, spread of invasive/alien species, and provide access for predators and incompatible recreation activities (Gunnison Sage-grouse Rangewide Steering Committee 2005). New and expanded highways, roads, and rail sidings have been built to service energy development, ranches, and residential properties throughout the Resource Area.




Figure 8. Location of power lines in the Resource Area. Data from Connelly et al. (2004), obtained from <http://sagemap.wr.usgs.gov/index.asp>.

### **C. Drought and Weather**

Long periods of below average precipitation, above average summer temperatures, above average snowfall, or below average winter temperatures, can adversely affect sage-grouse reproductive success and survival. In fact, prolonged drought during the 1930s and in the later part of the 20<sup>th</sup> century coincided with declines in sage-grouse populations throughout their range (Patterson 1952, Fischer 1994, Hanf et al. 1994). Extreme climatic conditions that occur during important life-cycle sequences have the potential to negatively affect food quality and abundance, as well as hiding cover (Hanf et al. 1994, Fischer 1996).

Sage-grouse can be very sensitive to fluctuations in annual moisture (Patterson 1952, Fischer 1994, Hanf et al. 1994). Sage-grouse summer diet, especially for chicks, is heavily dependent on insects and succulent plant growth. Sage-grouse population declines in some areas have been linked to years of low precipitation, most likely due to low nest success and/or poor chick survival (Hanf et al. 1994; Fischer 1996).

The Resource Area is part of the West Desert Basin. Precipitation reports from Basin monitoring stations indicate that the Resource Area experienced drought conditions from 2000 – 2004, during which time precipitation fell below the 30-year average (Figure 8). According to data collected at the writing of this Plan, 2006 was also a below average year.

Summer precipitation (April – September) appears to influence both sage-grouse habitat use and demographics on Deseret Ranch in Rich County, Utah. Sage-grouse use of lowland meadows increased in drier summers, and was negatively correlated with summer precipitation on Deseret Ranch (Danvir 2002). Data suggests populations were negatively affected by dry summers, as lek counts tended to increase following wetter summers while remaining stable or declining after drier summers (Danvir 2002).

Severe winter conditions can be a prominent factor in reducing grouse survival but there is no conclusive evidence to support this claim (Wallestad 1975; Beck 1977; Robertson 1991). Winter snow accumulations force birds to move to areas blown free of snow, or areas with sagebrush which extends above the snow (Eng and Schladweiler 1972; Wallestad 1975; Beck 1977; Hupp and Braun 1989; Robertson 1991). Bird loss can be significant in especially harsh winters, especially if the aforementioned conditions exist. The winter of 1983-84 was particularly severe, bringing extreme cold and heavy snow in Utah (and many parts of the western United States) for an extended period. It is believed that grouse populations declined dramatically during this winter. A far less severe, but still harsh, winter occurred in 1992-93. However, the impact of this winter on grouse populations in the WDARM area is not well documented.

Poor weather conditions in the spring are also suspected of influencing sage-grouse production (Connelly et al. 2000). Mild winters followed by relatively wet springs can increase production (Wallestad 1975, Autenrieth 1981) by promoting good insect and forb production. In contrast, severe spring weather (cold temperature combined with rain and wind) that coincides with hatching can decrease production as chicks die of exposure (Wallestad 1975).

Ongoing sage-grouse research will likely help to determine any potential effects of weather and drought on sage-grouse populations in the Resource Area.




Figure 8. Average precipitation for the Weber/Ogden River Basin 1995-2005 and the 30-year average (Utah Division of Water Resources 2006).

## **D. Hunting**

Connelly et al. (2000) maintain that most grouse populations can sustain controlled hunting seasons, but caution that grouse have the lowest reproductive potential of the upland game birds, that small populations (<100 male grouse counted during spring lek counts) are highly vulnerable, and that harvest rates should not exceed 10% of the fall population. Connelly et al. (2003) found that populations that are not exposed to hunting, recovered faster than populations receiving light to moderate hunting pressure. They recommend that grouse hunting seasons be conservative and account for population trend and habitat quality (Connelly et al. 2003a).

Field bag check data has been collected on sage-grouse in Utah since the 1960s and analysis of collected sage-grouse wings has been conducted since 1973. Grouse hunting pressure and harvest is obtained by two principal methods. The first method is to collect data from hunter questionnaires or telephone surveys. Harvest is also measured by collections of wings harvested by hunters.

Wing data also provides important information on other population parameters, including an estimate of annual chick production, assuming harvest represents the population. In the late 1980s and early 1990s, a drop in juvenile grouse per 100 hens was likely the result of drought conditions throughout much of Utah. During these years chick production was well below the recommended level of 225 juvenile grouse per 100 hens in the fall population.

Sage-grouse have been hunted legally in Utah since 1951. In 1988, Juab County was closed to sage-grouse hunting and Tooele County was closed in 1990. Closures were due to population declines (based on annual lek counts). Despite the closure of the sage-grouse season, UDWR hunter surveys continued to report sage-grouse hunters in the area, and hunters continue to report some sage-grouse hunting success (Tables 4 and 5). Hunter success information is obtained by the UDWR by mail-in surveys. Information for closed seasons, including Tooele and Juab Counties, may be an artifact from incorrect reporting by hunters, or is an indication of illegal activity. As it appears likely that information collected in some years is incorrect, it should be interpreted cautiously (D. Mitchell, UDWR, personal communication). In recent years, sage-grouse poaching has been suspected by biologists working in the field (J. Robinson, USU, personal communication) although this is not reflected in harvest reports from 2003-2005 (UDWR, unpublished data).

Table 4. Sage-grouse hunter harvest information for Tooele County, Utah 1984-2003 (UDWR 1984-2000, UDWR 2003). The County was closed to sage-grouse hunting in 1990.

Year	Hunter-days Afield	Birds Bagged	Birds per Hunter-day
2005	-	-	-
2004	-	-	-
2003	-	-	-
2002	-	-	-
2001	136	233	58
2000	-	-	-
1999	96	17	0.20
1998	0	0	0
1997	249	124	0.50
1996	185	92	0.50
1995	194	194	1.00
1994	210	386	1.83
1993	513	551	1.07
1992	355	231	0.65
1991	347	121	0.35
1990	267	308	1.15
1989	767	311	0.41
1988	985	267	0.27
1987	490	539	1.10
1986	267	412	1.54
1985	278	64	0.23
1984	304	162	0.53

Table 5. Sage-grouse hunter harvest information for Juab County, Utah 1984-2003 (UDWR 1984-2000, UDWR 2003). The County was closed to sage-grouse hunting in 1988.

Year	Hunter-days Afield	Birds Bagged	Birds per Hunter-day
2005	-	-	-
2004	-	-	-
2003	-	-	-
2002	-	-	-
2001	-	-	-
2000	-	-	-
1999	207	500	2.42
1998	93	93	1.00
1997	0	0	0
1996	69	278	4.00
1995	174	330	1.89
1994	123	333	2.71
1993	171	38	0.22
1992	106	17	0.17
1991	226	121	0.54
1990	308	61	0.20
1989	186	41	0.22
1988	133	116	0.88
1987	122	73	0.60
1986	82	247	3.00
1985	85	342	4.00
1984	121	40	0.33

## E. Fire

Across the Intermountain west, fire suppression is believed to have caused sagebrush stands to increase in canopy cover and density with a resulting reduction or loss of herbaceous understory species in many areas. Sagebrush stands have become more even-aged and less productive across large areas of sage-grouse habitat. Fires that do start, tend to burn greater acreage and at higher intensity due to the increased amount of fuel available to the fire.

The effects of any particular fire event depend on several characteristics of the local area including dominant sagebrush species, aridity, soils, topography, and disturbance (Bunting et al. 1987, Miller and Eddleman 2000). Other threats such as invasive/alien species (e.g. cheatgrass, *Bromus tectorum*), livestock grazing, and agricultural cultivation, are now present in sagebrush biomes, and contribute to the frequency, intensity, and duration of fire disturbances.

Fire, in general, is not detrimental to sage-grouse. In fact, sage-grouse have been observed to use burned areas so long as suitable cover and food are present during the time of use (Slater 2003). However, two altered fire regimes have emerged as being potentially incompatible with habitat management for sage-grouse populations. In the first, invasion of cheatgrass has increased the frequency of fire disturbances, which has the potential of changing sagebrush-steppe plant communities into grasslands (Miller and Eddleman 2000, Connelly et al. 2000). In the second, the occurrence of fire suppression has prevented the regular setback of succession and promoted the advancement of pinyon-juniper stands (Burkhardt and Tisdale 1969, Young and Evans 1981, Miller and Rose 1995, Miller et al. 2000). In these areas, there is the potential for sagebrush seed sources to be lost; reducing the likelihood that sagebrush could become reestablished after an eventual fire disturbance.

In the Resource Area, fire planning and management fall under the purview of land management agencies like the BLM, USFS, and local governments. The USFS Uinta National Forest operates according to a Fire Management Plan that is currently under revision. According to the draft 2005 Fire Management Plan (USFS 2005), the USFS fire management goals are to protect human life, both the public and firefighters, protect human communities, their infrastructure, and the natural resources on which they depend, and protect other property and improvements. The plan calls for the use of prescribed fire, wildfires, mechanical fuels reduction, and other available techniques to achieve these goals. The BLM is currently revising their fire plan (BLM 2005b).

Cheatgrass and pinyon-juniper encroachment appears to be pervasive and increasing in the Resource Area and may impact the way that fire management occurs in the Resource Area. Fire management by the BLM and the USFS is done in close cooperation with the UDWR who often provides a seed mix for post-burn rehabilitation. Fire planning is done carefully and cautiously in the Resource Area.

## F. Livestock Grazing

Livestock grazing is an important use of sage-grouse habitat in the Resource Area and throughout the range of sage-grouse in the West. The impacts of livestock grazing on sage-grouse are not clear, yet they are often contentious and controversial, perhaps more so than any other issue. Published literature and opinions run the gamut from completely compensatory or beneficial influence on one side, to incompatible, harmful practices that should be eliminated (Connelly et al. 2004). Due to the controversy, and following a thorough review on the subject by Rowland (2004), we have chosen to follow the lead of the Gunnison Sage-grouse Rangewide Steering Committee (2005), and simply provide several quotes from Rowland's publication. In addition, the potential impacts of livestock grazing on sage-grouse are covered extensively in Connelly et al. (2004).

### *Impacts to Sage-grouse Habitat*

Rowland (2004:17-19) summarized studies that suggest livestock grazing has a negative impact on sage-grouse habitat:

“Beck and Mitchell (2000) summarized potential effects of livestock grazing on sage-grouse habitats, and cited only four references that provide empirical evidence of direct negative effects of livestock grazing on sage-grouse, as follows. Of 161 nests examined in Utah, two were trampled by livestock (one sheep, one cattle) and five were deserted due to disturbance by livestock (Rasmussen and Griner 1938). In Nevada, sage-grouse habitat in wet meadows was degraded through overgrazing by domestic livestock and altered system hydrology (Oakleaf 1971, Klebenow 1985; as reported by Beck and Mitchell 2000). Klebenow (1982) examined sage-grouse habitat use in relation to grazing at the Sheldon NWR in Nevada, where sheep and cattle had grazed for >130 yr. Dominant sagebrush species at the refuge were low sagebrush, mountain big sagebrush, and Wyoming big sagebrush. Grasses included Sandberg and Cusick's bluegrass (*Poa secunda* and *P. cusickii*, respectively) in wet meadows, and Sandberg bluegrass and mat muhly (*Muhlenbergia richardsonis*) in dry meadows. A rest-rotation system was implemented for cattle grazing in 1980 over the majority of the refuge, where season-long grazing had occurred historically; a smaller portion had previously been managed under deferred rotation. Meadows heavily grazed by livestock (e.g., with few forbs and grasses and dense shrubs present) were avoided by sage-grouse, with the exception of use for free water when available (Klebenow 1982). (No explicit definitions were provided for light versus moderate or heavy grazing.)”

Rowland (2004:17-19) also noted cases where livestock grazing was reported to have had a positive effect:

“Some positive effects of livestock grazing were noted. When cattle were introduced into a meadow with residual grass, sage-grouse initially preferred the grazed openings, which had an effective cover height (sensu Robel et al. 1970) of 5 to 15 cm, compared to 30 to 50 cm in the lightly grazed surrounding areas. Grouse avoided dense, ungrazed basin wild rye meadows but were observed in adjacent wild rye that was grazed. One 40-ha meadow that was lightly grazed by cattle (41 yearling heifers, 60 days in June- August) was used

throughout the summer by sage-grouse and had more sage-grouse (100) than any other meadow on the refuge. Effective cover height in the meadow did not decrease below 5 cm during the summer.”

#### *Impacts on Sage-grouse Behavior and Demographics*

Studies that focused on sage-grouse behavior and demographic parameter response to grazing reported mixed impacts (Rowland 2004:17-19):

“Danvir (2002) reported two instances of nest abandonment related to livestock grazing in northern Utah during 7 yr of observations; one was caused by cattle, the other by sheep. Sage-grouse behavior on leks did not appear to be altered by the presence of cattle grazing (Danvir 2002). Sheep grazing in Idaho did not appear to disrupt use of leks by sage-grouse (Hulet 1983). Autenrieth (1981), however, cautioned against grazing sheep in sage-grouse winter habitat. He also suggested that livestock use of meadows occupied by sage-grouse, as well as livestock drives in sage-grouse habitat, could be detrimental to sage-grouse. In Wyoming, nesting densities of sage-grouse were considerably lower (10 nests/100 ha) in areas heavily grazed by domestic sheep compared to adjacent sites with moderate grazing (28 nests/100 ha) (Patterson 1952). Nest desertion caused by migrant bands of sheep also was documented (Patterson 1952). Heath et al. (1998) compared sage-grouse nesting and breeding success at three ranches with different grazing operations and levels of predator control in Wyoming. They found that, despite heavier livestock use (removal of >50% of annual herbaceous production, and grazing by both sheep and cattle) and long-term predator control on one ranch, nesting and breeding success of sage-grouse did not differ substantially among the three sites. Chick survival to 21 days was, however, greater on the ranch with lighter grazing, suggesting that predator control did not fully compensate for the greater reductions in herbaceous production (Heath et al. 1998). Further, hens were documented leaving the more heavily grazed ranch to nest elsewhere but returning to that ranch to rear broods (Heath et al. 1998). In a similar study, Holloran (1999) examined sage-grouse habitat use and productivity in relation to grazing management strategies at four ranches in southeastern Wyoming. He found no differences in nest success, brood survival, or numbers of chicks fledged among the ranches. Some differences in habitat use by sage-grouse were found among the ranches; however, these could not be ascribed to differences in grazing pressure, but were ascribed to differences in soil types and precipitation patterns (Holloran 1999). Above-average precipitation during the study, however, may have obscured any potential differences in habitat suitability for sage-grouse among sites. Neither of these studies employed control sites or replication.”

#### *Recommendations*

In her extensive literature review, Rowland (2004:11) summarized recommendations found in the literature related to timing of grazing and reduction of impact to riparian areas used during brood-rearing. In addition, Rowland (2004: 24) made her own recommendations:

“Timing of grazing greatly influences the effects of livestock grazing in meadows and riparian areas. These sites are particularly vulnerable in late summer when excessive grazing and browsing may damage riparian shrubs, reduce the yield and availability of

succulent herbs (Kovalchik and Elmore 1992), and cause deterioration of riparian function over time (Klebenow 1985). However, moderate utilization by livestock in spring, early summer, or winter is sustainable in non-degraded meadow and riparian areas within sagebrush habitat (Shaw 1992, Clary et al. 1996, Mosley et al. 1997). Moderate use equates to a 10-cm residual stubble height for most grasses and sedges and 5-cm for Kentucky bluegrass (Mosley et al. 1997, Clary and Leininger 2000). Shrub utilization should not exceed 50-60% during the growing season, and at least 50% protective ground cover (i.e., plant basal area + mulch + rocks + gravel) should remain after grazing (Mosley et al. 1997). While hydrophytic shrubs may not directly serve as sage-grouse habitat, they do impact the stability of riparian and meadow habitats important to sage-grouse (Winward 2000). The length of time livestock have access to meadows may be more important than the level of utilization; it has been suggested that livestock access be limited to 3 weeks (Mosley et al. 1997). In riparian and meadow habitat degraded by heavy livestock utilization, rest from grazing may be necessary for recovery (Clary and Webster 1989).”

“Manage livestock grazing through stocking rates and season of use on all seasonal ranges of sage-grouse to avoid habitat degradation (Paige and Ritter 1999, Beck and Mitchell 2000, Wisdom et al. 2000), especially on recently disturbed sites, such as those sprayed or burned (Braun et al. 1977). In nesting and brood-rearing habitats, ensure that grazing does not reduce herbaceous understory cover below levels that serve as a deterrent to potential predators of eggs and chicks (Connelly et al. 2000, Hockett 2002). Healthy native understories also support insects and forbs that are important in diets of pre-laying hens and chicks (Johnson and Boyce 1990, Barnett and Crawford 1994, Drut et al. 1994b). Riparian areas and wet meadows used for brood rearing are especially sensitive to grazing by livestock; in these habitats, removal of livestock before the nesting season may be prudent (Beck and Mitchell 2000, Hockett 2002).”

### *Conclusions*

Livestock grazing is an important use of sagebrush rangelands in the Resource Area. Although some incompatible grazing likely occurs within the Resource Area, the majority of livestock operations appear to be coexisting with sage-grouse. However, no studies have been conducted in the Resource Area to specifically address the issue of grazing impacts on sage-grouse and this is an area that may warrant future research.

## **G. OHV Recreation**

The effects of off-highway-vehicle (OHV) recreation and other forms of recreation (snowmobiles, bird watching, etc.) on sage-grouse behavior and populations are poorly understood. Impacts of recreational activities are likely to be of two forms: disturbance of individuals and alteration of habitat.

Recreational activities, specifically OHV recreation, likely has the potential to impact individual birds, or flocks of birds, by flushing them from breeding grounds, nests, roost sites, or foraging areas, depending on the season in question. Noise associated with OHV recreation is likely the primary cause of disturbance to individual or flock behavior. Disturbance during nesting season may result in nest abandonment or failure. Disturbance during any time of year may increase the vulnerability of sage-grouse to predators. OHV recreation, and other forms of recreation, may also trample plants, disturb soils, and otherwise alter and degrade habitat. In many instances, specific areas are designated for use of OHVs. When confined to specific use areas, impacts are likely to be reduced.

OHV recreation is relatively common in the Resource Area. Although specific impacts to sage-grouse populations are unknown, there is some concern that OHV recreation impacts sage-grouse use areas. Tooele County is actively planning for expansion of OHV trail systems in and around the town of Tooele (Figure 9). We cannot predict what impacts additional trail systems may have on sage-grouse in the Resource Area; monitoring of this threat is recommended. In addition, we recommend investigating, in a GIS system, where proposed trails may impact crucial sage-grouse habitats as identified by the UDWR and/or on-going research efforts.




Figure 9. Existing and proposed trails in Tooele Valley (Tooele County 2005).

## H. Invasive/Noxious Weeds

The Utah Department of Agriculture (Section 4-17-2) defines noxious weeds as, "...any plant the commissioner determines to be especially injurious to public health, crops, livestock, land, or other property." Under the Utah Noxious Weed Act (4-17-10) county weed departments are charged to, "...develop, implement, and pursue an effective program for the control and containment of noxious weeds on all lands under their control or jurisdiction, including highways, roadways, rights-of-way, easements, game management areas, and state parks and recreation areas."

Russian knapweed (*Centaurea repens*), dyers woad (*Isatis tinctoria* L), and several other species of thistle, grasses, and knapweed are listed on the Utah Noxious Weed List (Section 4-17-3, Utah Noxious Weed Act). Although cheatgrass is not listed there, nor is it included in individual county lists for Tooele and Juab Counties, this invasive plant species is known to be established in the Resource Area.

Invasive species affect the species composition, nutrient cycling, and physical structure of sagebrush systems. Invasive species also influence the natural function of sagebrush systems, especially their ability to recover from fire. These impacts often culminate in an alteration of wildlife species diversity and abundance in affected systems.

Cheatgrass is an annual grass native to Russia and parts of northern Europe. When it invades sagebrush communities, cheatgrass is known to increase fire frequency and has the potential to convert sagebrush communities to annual grass rangelands. Cheatgrass has also been reported to encourage establishment of other invasive species (Grahame and Sisk 2002).

Noxious weeds have been recognized within the Resource Area as a serious problem by County Weed Control departments, BLM, and USFS. County weed control departments maintain records of the location, extent, and severity of weed establishment, and actively work to control the spread and establishment of weeds in their respective counties. In January 1996, the BLM published Partners Against Weeds, (PAW) an action plan for the Weed Management program in the Bureau. The PAW plan lists seven goals, the first being to develop a prevention and early detection program. The PAW recommends developing and enforcing a policy to "ensure seeds, seed mixtures, hays, grains and straws are free of weed seed" as a prevention and detection strategy. Utah's BLM Resource Advisory Council (RAC) developed a guideline requiring certified weed free forage to be used on BLM lands by anyone having the need to take forage with them when using BLM public lands. The guidelines were approved by both the Utah State Director and the Secretary of the Interior in 1997. Beginning in November 1998, users of BLM administered land in Utah are required to use only certified noxious-weed-free hay, straw, or mulch. Approved products for livestock feed on public lands include pellets, hay cubes, processed grains and certified hay, straw or mulch, all of which are normally available at many feed stores and producers in Utah. The USFS is also committed to a campaign against the spread of invasive species. Working with agency and local government partners, the USFS aims to create Cooperative Weed Management Area (CWMA) Participative Agreements for all USFS lands.

## I. Parasitism and Disease

Several bacterial and parasitic diseases may affect sage-grouse to varying degrees. Sage-grouse have long co-existed with a range of pathogens and many produce no or few, ill effects in individuals and populations. Large-scale (i.e. rangewide or statewide) impacts to sage-grouse have not been reported. Below, we discuss a few of the pathogens that appear to be most likely to impact sage-grouse populations (Connelly et al. 2004).

### *West Nile Virus*

West Nile virus (WNV) is an arbovirus, or arthropod-borne virus, of the flavivirus family, which also includes Dengue and Yellow Fever. WNV is one of many mosquito-borne viral infections. Mosquitoes of the *Culex* family primarily transmit West Nile Virus during normal blood feeding. Some species in this family feed primarily on birds, and birds act as reservoirs or amplifying hosts of the virus. Although many species of birds are known to contract WNV, species in the Corvid family (crows, ravens, and jays) are more susceptible to the disease and are therefore useful geographic detectors of WNV. Mammals, including humans and horses, are considered incidental hosts and are therefore viral 'dead ends.' Humans are most likely to acquire WNV from an infected mosquito. Other mammals, such as horses, do not maintain a sufficiently high level of the virus in the bloodstream to transmit the virus to humans.

WNV was first detected in the Western Hemisphere in 1999, and has since rapidly spread across the North American continent into all 48 continental states, seven Canadian provinces, and throughout Mexico. In addition, WNV activity has been detected in Puerto Rico, the Dominican Republic, Jamaica, Guadeloupe, and El Salvador.

In 2003, several cases of WNV were confirmed in sage-grouse in Wyoming (nineteen birds), Montana (three birds), and Alberta, Canada (five birds). In that same year, WNV was detected in chickens in Price, Utah. In 2004, sage-grouse in Wyoming, Montana, Colorado, and California tested positive for the virus. In 2005, the virus was confirmed in a dead sage-grouse in the Uintah Basin, approximately 156 miles to the east of the Resource Area. WNV was also detected in a prairie falcon in Carbon County, to the southeast of the Resource Area. A limited percentage of sage-grouse appear to be capable of developing immunity to the virus (Cornish, unpublished data) and infection appears to be almost always fatal within 24–48 hours.

### *Macro-parasites*

Coccidiosis—Coccidiosis is an intestinal disease caused by one or more species of the protozoan genus *Emeria* (Jolly 1982), which include *E. angusta*, *E. centroceri*, and *E. pattersoni*. Infection results in diarrhea caused by damage to the mucosal lining of the digestive tract. The disease is transmitted through consumption of contaminated feces. Coccidiosis is the most well known of all diseases infecting sage-grouse (Connelly et al. 2004). In Wyoming, Colorado, and Idaho from 1932–1953 this disease resulted in significant losses of young sage-grouse (Hones and Post 1968), however no cases have been documented since the 1960s (Connelly et al. 2004). Cases were typically reported in areas where large numbers of birds are concentrated. The concentration led to contamination of and spread via water and food sources. Connelly et al. (2004) speculated that this disease lacks prevalence in recent years because sage-grouse density

has decreased. No cases of Coccidiosis are known from within the Resource Area, however this does not imply that the condition does not exist or have the potential to exist. Specifically, drought conditions that result in a decrease in water sources may potentially increase sage-grouse concentrations in localized areas, thereby increasing the potential for impacts from this infection.

Tapeworms—Sage-grouse are the only known host of the cestode tapeworm, *Raillietina centroceri* (Honest 1982). There is little consensus on the impact *R. centroceri* may have on sage-grouse populations. The Canadian Sage Grouse Recovery Strategy indicates that this infection may be a largely overlooked cause of mortality. Honest (1982), suggested that there was a synergy between host and parasite with little negative impacts to sage-grouse. The parasite does not affect the quality of sage-grouse meat and there are no documented cases of *R. centroceri* in the Resource Area. This does not imply that this infection does not impact sage-grouse therein, however.

Filarid Worms—A filarial nematode, *Ornithofilaria tuvensis*, which utilizes the connective tissue between skin and breast muscle in sage-grouse, appears to prevent flight in infected birds (Hepworth 1962). This infection is rare but appears to have significant impacts. This infection is not known to occur in the Resource Area, although it may yet exist, undetected.

Avian Malaria—Avian malaria, caused by the protozoan *Plasmodium pediocetti*, is known to infect wild sage-grouse but is considered rare. Although this infection does not have a profound impact on sage-grouse populations, it does cause birds to reduce activity during morning hours and may affect courtship and breeding of strutting males (Boyce 1990, Johnson and Boyce 1991). Biting flies (Friend and Franson 1999) transmit this disease.

### Conclusions

We currently consider WNV to be the disease/parasite with greatest potential to impact sage-grouse populations in the Resource Area. As previously mentioned, in 2005 a dead sage-grouse was found in the Uintah Basin 156 straight-line miles to the east of the Resource Area. The sage-grouse was infected with WNV; however there is no interaction between birds in this area. At the time of writing, WNV has been detected in one person, one horse, eight chickens, and several mosquitoes in the Resource Area (Utah Department of Health 2006). Continued monitoring of radio-collared birds (Robinson, unpublished data) will help to detect this pathogen in the population early.

Other diseases discussed in this section may have an effect on sage-grouse but only one has been observed in the Resource Area and, therefore, is considered a limited potential threat.

## J. Predation

Sage-grouse occupy an important place in the food web in sagebrush environments and are preyed upon by a wide variety of terrestrial and avian predators. Numerous predators have been documented preying upon differing ages of sage grouse and/or their nests. Documented nest predators include weasel (*Mustela* spp.), badger (*Taxidea taxus*), elk (*Cervus elaphus*), coyote (*Canis latrans*), common raven (*Corvus corax*), American crow (*Corvus brachyrhynchos*), red fox (*Vulpes vulpes*), striped skunk (*Mephitis mephitis*), black-billed magpie (*Pica pica*), and various species of snakes (Family *Leptotyphlops*; Batterson and Morse 1948, Patterson 1952, Nelson 1955, Autenrieth 1981, Hanf et al. 1994, Young 1994, DeLong et al. 1995, Sveum 1995). Numerous species have also been documented killing and/or consuming adult sage-grouse and include golden eagle (*Aquila chrysaetos*), Cooper's (*Accipiter cooperii*), ferruginous (*Buteo regalis*), red-tailed (*Buteo jamaicensis*), and Swainson's hawks (*Buteo swainsoni*), Northern goshawks (*Accipiter gentilis*), coyote, red fox, and bobcat (*Felis rufus*; Girard 1937, Rasmussen and Griner 1938, Batterson and Morse 1948, Nelson 1955, Rogers 1964, Beck 1977, Dunkle 1977, Autenrieth 1981). Numerous predator species, many of which are listed above, have been documented killing juvenile sage-grouse. Because of the small size of young sage-grouse, additional predators have been documented and include American kestrels (*Falco sparverius*), merlin (*Falco columbarius*), Northern harrier (*Circus cyaneus*), common raven, and weasel (Girard 1937, Patterson 1952, Nelson 1955, Rogers 1964, Autenrieth 1981).

Predation is the end result for the vast majority of sage-grouse throughout their range, both historically and presently (Bergerud 1988). Schroeder and Baydack (2001:26) suggest that predation has the potential to affect the annual life cycle of sage-grouse in three primary ways:

1. Success of nests
2. Survival of juveniles during the first few weeks after hatch
3. Annual survival of breeding-age birds

Peterson and Silvy (1996) conclude that the relative importance of predation on the viability of sage-grouse populations is relatively unknown and warrants additional study.

Nest success varies by year, area, population density, and/or management strategy (Connelly et al. 1998, Schroeder et al. 1999). Connelly et al. (2000) suggested that several studies on nest success have illustrated success >40% and that nest predation does not appear to be a problem across the range of the grouse. In contrast, Gregg (1991) and Gregg et al. (1994) suggested that nest predation may be limiting grouse numbers in Oregon. Red foxes and common ravens have been implicated in affecting nest success and the annual survival of breeding age birds in the Strawberry Valley area of Utah (Bunnell et al. 2000). Researchers suggest that the advancing population of a nonnative predator, red fox, is responsible for preying upon a large portion of the population in that area (Flinders 1999). In artificial nest studies conducted in Strawberry Valley, ravens depredated 98% of artificial nests within 48 hours of their placement; remote cameras were used to verify the identity of artificial nest predators (Baxter and Flinders, unpublished).

### *History of Predator Management in Utah*

*The following sections on the history of predator management and effects of predator management on sage-grouse populations were written, in conjunction with WDARM, for the*

*Plan by representatives from USDA-WS. Wildlife Services has been managing predator populations and collecting data on predator population trends in the state for several decades. They also have an extensive body of personal knowledge about predator population management and the impacts of various management practices.*

Understanding the impact of predation on sage-grouse is difficult, as the primary effects (the number of sage-grouse killed by predators) is affected by habitat variables, variables associated with the predator population, and variables within the sage-grouse population itself. Secondary effects of predation exist and are indicated when habitat choices are dictated by the risk of predation. What we currently know about habitat needs of sage-grouse is developed from studies of core sage-grouse range. However, if predation or the risk of predation is effecting habitat selection, then otherwise good habitat is made unavailable to sage-grouse. To better understand the role predation management may have played historically, it is important to examine records of the past.

Predator management in Utah began in the late 1800s with territorial bounties followed by a federal appropriation in 1917. The original purpose for the federal program was the suppression of rabies. The program has gone through several changes involving both State and Federal agencies. The U.S. Biological Survey managed predator control in early years and developed the structure that was later used by the USFWS which is to have supervised men in designated Wildlife Service districts. From 1936 to 1986, the USFWS managed the program as Animal Damage Control. In 1986, it moved to the United States Department of Agriculture under the Animal and Plant Health Inspection Service, and in 1996 was renamed again as Wildlife Services (USDA-WS).

Correlations exist between livestock inventories and the intensity of predator management efforts. Domestic sheep numbers are on record (Utah Agricultural Statistics) at a high of 2.7 million in 1931. Breeding sheep inventories as of January 1, 2003 were reported at 290,000 head, or 10.7% of the maximum number. Sheep numbers varied substantially from year to year in some cases, and from decade to decade since the early 1900s. Toxicants were used extensively in the early years when sheep numbers were high. Additionally, predator management in the early years involved many trappers, setting and tending steel traps statewide. As many as 132 men were hired (1936) to set traps and apply baits. Figure 9 shows the recorded take of coyotes from the predator control program between 1917 and 2004. These data do not include poisoned coyotes, which were not found but estimated at seven to ten coyotes for each dead one found.




Figure 9. USDA-WS reported coyote take in Utah, 1917-2004. Data reported by USDA-WS.

Strychnine and thallium treated single-lethal-dose (SLD) baits were the main toxicants used between 1920 and 1950. Compound 1080 was developed around 1945, first as a rodenticide and later as a predicide used in large bait stations. USDA-WS records indicate that 1443 bait stations were applied in 1969, covering 54% of the townships in the State. Bait placement from 1950 to 1972 were large bait stations while SLD baits were used prior to that. From about 1950 to 1972, Compound 1080 became the main force in controlling coyotes in all districts of Utah by the government and by private individuals. The low government take of coyotes during this period indicates that coyote populations were suppressed by bait station use. Toxicants were banned in federal programs in 1972, and current policies allow only two very selective toxicants for limited use.

Early predation control also extended to ravens, crows, eagles and magpies. Records indicate that single baits were applied around ‘draw stations’ to target birds. Records also note that UDWR personnel targeted specific areas for bird suppression not treated by the federal program.

It is difficult to assess the poison years in terms of population suppression of species such as ravens, coyotes, and even red foxes. Individual species records are presented below:

Red Fox—While some early records of red foxes exist, red foxes are believed to have been virtually absent on the landscape before the 1970s. Red fox do not exist in government records before 1972, and have increased since then. Red foxes may have been successfully suppressed by rabies or by bait station use, or both. Figure 10 shows red fox take from 1972 to 2004.




Figure 10. USDA-WS reported fox take in Utah, 1972-2004. Data reported by USDA-WS.

It is interesting to observe that the statewide increase in red fox abundance has occurred during the lowest period of coyote control. Sargeant et al. (1984, 1987) theorizes that protection for coyotes would allow coyote populations to increase, thus suppressing red fox populations. This has not happened on a landscape basis in Utah.

An argument may be made that red fox in Utah are an invasive species, based on historical data. Red foxes were historically divided into two species, *Vulpes vulpes* in the Old World and *V. fulva* in the New World, but today they are considered to be one species in the U.S. (*V. vulpes*). Churcher (1959) reviewed twelve subspecies of red foxes in North America with nine subspecies currently recognized in Canada. Churcher (1959) suggested that the red fox was introduced from Europe to the southern colonies around 1790.

Following the introductions, there was confusion as to which populations were expanding. Audubon and Bachman (cited in Churcher 1959) believed that Pennsylvania was the southern limit of the red fox's range in 1750, and documented a range extension southwards to Georgia by 1850. Leopold (1933) reported the expansion of red fox in Wisconsin, which was displacing the grey fox, while Godin (1977) reported *V. fulva* had established itself by 1850 and was displacing the gray fox to some degree along the southeastern seaboard. Godin also speculated that the introduced foxes might have interbred with a scarce population of indigenous red foxes, but historical accounts do not support this. Churcher (1959) concluded from the available evidence that the red fox was native to North America north of 40-45°N but was scarce or absent in the hardwood forests where gray foxes were common. Churcher (1973) suggested that the 'original' habitat was the northern mixed hardwood and softwood forest zones. He also observed that the red fox might have been found in the hardwoods to the south and the tundra to the north. Gilmore (1946) believed that red foxes were absent from Pennsylvania during aboriginal times and concluded that they did not range into the mideastern United States. Rhoads (1903 cited in Churcher 1959) stated, "in earlier colonial times the red fox was unknown in the austral zone (southern states)."

Archaeological evidence from Ontario, Canada (Peterson et al. 1953) has shown that the red fox was present in the Midland area prior to introductions during the decade 1639–1649 and that it was present earlier in the Oxford and Middlesex counties of southern Ontario, Canada. Sites farther south did not have red foxes (Gilmore 1946).

Once the red fox began to spread south and west from northeastern U.S. after its introduction from Europe, it expanded its range to include the prairies of the mid west and continued to expand west to Colorado and Utah. It has reached the Utah–Nevada line, and seems likely to invade Nevada as well.

Striped Skunk—Historically, what may be significant is the relatively few skunks found in Utah. Figure 11 shows skunk take by USDA-WS in Utah from 1917-2004. Periodic rabies eruptions suppressed skunk populations in the early years of the century. As an example, in 1918 with 51 full-time personnel setting traps, only ten skunks were removed statewide. In the 1920s, following years of SLD bait placements, skunk take in the program increased to above 100 annually, but then declined to none in 1933, 12 in 1934, 35 in 1935, and up to 98 in 1936. The cycle of skunk removal probably reflects the population level effect of rabies in skunks. The last skunk rabies incident in USDA-WS records, occurred in 1972 in Davis County, with a

countywide control program initiated as a result.




Figure 11. USDA-W.S. reported skunk take in Utah, 1917-2004. Data reported by USDA-W.S.

Raven and Magpie—Breeding bird survey results indicate a 300% increase in raven numbers from 1968 to the present. While most biologists believe the increase is due to more favorable conditions and anthropogenic food sources, the increase in populations also follows the reduction in use of poisons, likely keeping their numbers low.

Magpies were targeted along with ravens at draw stations with smaller baits placed especially for birds. USDA-W.S. records show that UDWR personnel placed baits to target birds in areas where federal poison programs were not active.

Coyote—Intensive coyote control prior to 1972 suppressed the coyote population. Since that time, the design of the predation management program has been to reduce coyote damage while not influencing populations. Analyses by Connolly and Longhurst (1975) and Pitt et al. (2002) indicate that the current level of exploitation does not affect coyote populations. It seems likely that coyote populations have never been higher than in modern times.

#### *Impacts of Predation on Sage-grouse*

Given that predators and nest predators are abundant and many are present in all time high numbers, impacts to sage-grouse may take one of two forms. Sage-grouse may be killed directly by predators (primary effects), and most mortality of sage-grouse is predation. Direct predation has negative population effects when it exceeds recruitment.

Secondary effects of predation include biological effects that are the result of behavioral changes

in sage-grouse. These behavioral changes result from the risk of predation and may take the form of lower fecundity, longer dispersals, use of sub optimal habitat, nest abandonment, and a number of other behaviors, which may affect populations.

Autenrieth (1981) suggested that nest predation was likely the most important population constraint on sage grouse in his study area. Here, predation on adult birds does occur and may be significant in some cases. Presnall and Wood (1953) reported tracking a coyote approximately five miles to its den in northern Colorado, and finding evidence along the way that the coyote had killed three adult sage-grouse and destroyed a sage-grouse nest. Examination of the stomach contents from an adult female coyote removed the next day showed parts of an adult sage-grouse plus six newly-hatched sage-grouse chicks. The area around the den site was littered with sage-grouse bones and feathers. No other prey animal remains were found around the den, and it appeared that the pups had been raised largely upon sage-grouse. Till (1992) documented sage-grouse remains at 4 of the 30 coyote den sites examined during his study in south central Wyoming, but provided no indication of the relative abundance or distribution of sage-grouse in his study area. In northern Utah, researchers from Brigham Young University confirmed predation, primarily by red fox and coyote, as the cause of death for 13 of 21 radio-instrumented sage-grouse in the first year in their study area (Bunnell and Flinders 1999). Two additional instrumented birds could not be found, but were suspected to have been killed by predators, suggesting a 71% predation loss of instrumented birds. Additionally, eleven other sage-grouse were found dead in their study area, and all but one of these birds was killed by mammalian predators. USDA-WS is not aware of controlled studies conducted to determine if coyote and red fox control would actually result in significant benefits to grouse populations. However, the above studies indicate there may be some benefit to the removal of these predators in some situations.

In addition to primary predation effects secondary predation impacts likely exist in a number of populations. The risk of coyote predation may cause habitat abandonment or, through habitat choices, reduce fitness and make grouse more susceptible to other mortality. Coyote damage management may be indicated for populations not performing to habitat potential.

**Meso-predator Release**—As red fox have been implicated as primary predators of sage-grouse in many areas, the notion of some natural control of red fox by coyotes has been suggested. The idea that coyote removal may benefit red fox, and thus be a detriment to sage-grouse, has been offered by some as a need to limit coyote removal. The potential for an indirect effect on sage-grouse of coyote removal would take the form of a ‘meso-predator release,’ which is the increase in smaller mammalian carnivore species after larger carnivores have been reduced or eliminated. The meso-predator release theory allows that smaller predators are allowed to increase due to either a lack of predation, a release from competition, or both. Gehrt and Clark (2003) present an opposing view of meso-predator release and point out several weaknesses in the circumstantial evidence that has been used to suggest that meso-predator release occurs.

Sargeant et al. (1984) reported on the effects of red fox predation on breeding ducks. Their data were collected when coyote populations were presumably suppressed by widespread use of predicide, and he notes that at the time (1968–1973), “[c]oyote populations in most of the midcontinent area appear to be suppressed by man.” The authors noted an inverse relationship between red fox and coyote populations and speculated, “...protection of coyotes will result in

expansion of local or regional populations that in turn will cause reductions in fox populations.” They inferred that this would reduce predation on upland nesting ducks. Sargeant et al. (1987) reported on spatial relationships between coyotes and red foxes and showed that home ranges of fox families did not overlap the core centers of coyote home ranges on a North Dakota study site. Although none of their radio-collared foxes were killed by coyotes in their study, they hypothesized that red foxes tended to avoid coyote territories, presumably because of the fear of being killed by coyotes. Thus, they inferred that a red fox population would increase if the coyote population were reduced, because removal of territorial coyotes would create vacant coyote territories that could then become occupied by red foxes.

Still, the presence of coyotes does not completely displace red foxes. Voigt and Earle (1983) verified that red fox travel through coyote areas during dispersal, but did not establish there. They also reported, “...individual foxes and coyotes can occur in close proximity to each other along territory borders and when coyotes travel into fox areas.” They also noted that “fox-coyote range overlap near borders was similar to fox-fox range overlap near borders,” and that coyotes do not, “completely displace foxes over areas.” Gese et al. (1996) reported that coyotes tolerated red foxes when encountered about half of the time in Yellowstone National Park, although at times they were aggressive toward, and would sometimes kill, foxes.

Other studies suggest that coyote territories would not remain vacant for very long after the coyotes are removed. Gese (1998) noted that adjacent coyote packs adjusted territorial boundaries following social disruption in a neighboring pack, thus allowing for complete occupancy of the area despite removal of breeding coyotes. Blejwas et al. (2002) noted that a replacement pair of coyotes occupied a territory in approximately 43 days following the removal of the territorial pair. Williams et al. (2003) noted that temporal genetic variation in coyote populations experiencing high turnover (due to control) indicated that, “...localized removal did not negatively impact population size...” In Utah, USDA-WS removes a small percentage (2-4%) of the estimated coyote population, not enough, even at a small scale, to create the vacant territories that would theoretically allow red fox populations to increase substantially. Therefore, we believe it would be unlikely for USDA-WS coyote removal actions to lead to indirect increases in predation effects on grouse populations. To the contrary, where populations are not performing to the full potential of the habitat, predation management may be necessary as part of an applied management plan for sage-grouse.

**Predation Defense Mechanisms**—Sage-grouse have adapted to live, and have evolved with, many of these predators. Sage-grouse, and other ground nesting birds, have developed effective strategies for hiding from predators when they occupy habitat of sufficient quality. Schroeder et al. (1999) briefly describe some of those adaptations. The actual timing of the strutting display and/or the formation of leks may have evolved due to predation selective pressures (Patterson 1952, Hartzler 1972, Bergerud 1988, Phillips 1990). Sage-grouse also respond to predation by either crouching in dense vegetation or flying away from an attacking predator (Hartzler 1972, Ellis 1984). Female Greater Sage-grouse have also been documented defending their nests from ground squirrels (Schroeder 1997). Girard (1937) observed females attacking predators in the defense of their brood. In an attempt to lead potential predators away from nests or young chicks, females have been documented performing distraction displays. The distraction display includes dragging wings on the ground while moving erratically (Peterson 1980). In addition, a female will occasionally re-nest if her first nest is destroyed by predators early in the incubation

period (Patterson 1952, Eng 1963, Connelly et al. 1993, Schroeder 1997), although re-nesting rates for sage-grouse are relatively low (Connelly et al. 1993).

#### *Predator Control and Livestock Populations*

Predator control activities began in Utah in 1888 with Territorial Bounty laws, which continued into statehood and through the early 1900s. The inauguration of the government sponsored predator control program began in 1915 with small appropriations of funds used to hire a supervisor and eight men in designated areas where control was needed to protect livestock. Today this program is managed by USDA-WS.

Utah sheep numbers were at a record high of 2.7 million in 1931. Numbers varied from year to year in some cases and from decade to decade since the early 1900s. Today, approximately 265,000 sheep are grazed in Utah. Although sheep numbers are down, more cattle ranching operations exist in the state today compared to 1931.

Predator control for the protection of cattle replaced some of the reduction in control because of reduced sheep numbers. Improved methods of hunting with aircraft increased efficiency and effectiveness since the early 1970s, but poisons were used extensively in the early years when sheep numbers were high. Congress passed the Animal Damage Control Act of March 2, 1931. Records show that in 1936 up to 132 men were hired for predator control. Poison baits placed by men in the various field districts were more effective at controlling predator populations over a larger area than are currently worked today. Government trappers took a documented 16,719 predators in 1939, yet that figure doesn't reflect all of those which were poisoned. This amount was a record catch for any one fiscal year and shows more predators were taken in early years than records of today. The number of predators taken during this era not only exceeds the modern 'take,' but likely represents a larger percentage of the population of the day. Modern records (since 1972) show that on average, USDA-WS in Utah averages about 5,000 coyotes per year by using 25 field men and several fixed-wing aircraft along with contracted helicopter work. Another 5,000 coyotes (on average) are taken by private hunters and trappers annually in Utah.

Utah's coyote population today is near 100,000 based on studies by USDA-WS research personnel (Connolly 1996). Predator damage management today focuses on individuals causing damage, as opposed to population reductions (or eradication in the case of the wolf) of the past. Current control is practiced on less land mass, with more restrictions, and for the protection of fewer livestock than at any time in Utah history. Correspondingly, there are probably more coyotes alive today than at any time in Utah history.

Strychnine and thallium were the main poisons used in the early 1900s until the advent of Compound 1080 in 1945. Compound 1080 was first effectively used on rodents and later on predators. From about 1950 to 1972, Compound 1080 became the main method in controlling coyotes in all districts of Utah by the government and by private individuals. It is impossible to know precisely the effects it had on the coyote population, as population census were not conducted and the main objective of control was eradication.

It is reasonable to believe that Compound 1080 reduced coyote numbers considerably in large tracks of land that are no longer worked because current land-use practices prohibit coyote

control. Strychnine baits used for coyote control before 1972 (in conjunction with Compound 1080) likely controlled ravens and raptors, which fed on the baits. Compound 1080 is highly selective to canines but overused because there were no dosage restrictions or regulations in place.

It is difficult to assess the extent of population suppression for ravens, coyotes, and even red foxes during the poison years. Some red foxes were found in Utah in low numbers and at high elevations early in the Territorial history. However, most biologists believe the red fox in Utah today is an invasive species, which arrived in the 1970s. Ravens have increased in numbers from the 1970s likely due to more favorable conditions, including human food sources (landfills, etc.). The increase in the raven population also follows the reduction in use of poisons that could have kept their numbers low. Early records show raven predation on lambs in the 1950s and concern to control them.

The effects of reduced coyote control on sage-grouse are not well understood. The decline of sage-grouse occurred at the same time as coyote populations expanded. It could be concluded that the poison ban allowed coyotes, raptors, and ravens all to expand in population numbers and range. Protections were placed at this time on ravens and magpies in the form of removing bounties and adding laws that prohibited shooting and nest destruction. The reduction in sheep numbers added to the favorable habitats for predators and raptors by the increases in prey base, and improved meadows and riparian areas. Red foxes arrived at this time and expanded in numbers because of the more favorable environments as previously discussed.

Incidentally, sage-grouse could have also benefited from the high numbers of sheep concentrated in winter areas. Properly managed sheep grazing in the winter has the effect of rejuvenating sagebrush. As sheep numbers declined, sagebrush became decadent to some degree. Although natural cycles may have once occurred in sage-grouse populations, changes in the environment since the 1970s have caused a long term decline. Once a decline in sage-grouse numbers occurred, the increase in predator numbers, especially red fox and ravens, would be more detrimental to the grouse.

### *Conclusions*

On-going research has documented predation events on sage-grouse nests, juveniles, and adult birds (Appendix B). Many sage-grouse predators are known to occur in the Resource Area and USDA-WS does conduct predator control in the area related to livestock operations which is likely to influence predator-prey dynamics involving sage-grouse. Predation by both native and nonnative predators is considered a relatively serious threat to sage-grouse populations in the area. There is some concern that increasing populations of red foxes, other nonnative predators, and anthropogenic population inflation of common ravens in the Resource Area could be having a negative impact on sage-grouse populations. Additional research and investigation into this issue seems warranted.

## **K. Renewable and Non-renewable Energy Development**

There is little oil and gas activity in the Resource Area, as described below. Although mining has traditionally been an important resource in the Resource Area, impacts are generally restricted to higher elevation areas with rugged terrain where sage-grouse are not found.

Oil and gas facilities generally have a small footprint, usually a few acres or less. Each pad will often contain tanks and other equipment for a period of years. When the well is depleted, all facilities are removed and the pad is reclaimed. Some researchers believe the existence of these facilities suppresses sage-grouse use of the habitat for some distance beyond the actual footprint of the facility (Robel et al. 2004, Holloran et al. 2005). Compressor stations, active wells, and drilling rigs produce relatively loud and sustained noise that may interfere with sage-grouse, particularly during the breeding season (Crompton et al. 2006).

Since 1991, only four wells have been spudded (drilling commenced) in the Resource Area. The location of drill sites is depicted in Figure 13 and is provided to illustrate generally, where oil and gas impacts are located in the Resource Area.

Effective reclamation of oil and gas pads and other facilities, including the reestablishment of big sagebrush in some instances, is important for maintenance of sage-grouse habitat in these development areas. Although this can be challenging in drier portions of the Resource Area, reclaimed pad sites have been used by sage-grouse in some parts of Utah (B. Maxfield, UDWR, personal communication).




Figure 13. Location of drill sites in the WDARM Resource Area. Data obtained from [http://www.ogm.utah.gov/oilgas/MAP\\_SEARCH/map\\_search.htm](http://www.ogm.utah.gov/oilgas/MAP_SEARCH/map_search.htm)

## **L. Vegetation Management**

Vegetation management conducted in the past was a reflection of the priorities of the time, and on the mandates and policies of the federal government, when vegetation management was done on federal land. Because much of the land in the Resource Area is under federal management, this is an important consideration when evaluating past and current conditions. In the past, many vegetation treatments were conducted to increase forage for livestock.

Recently, vegetation management has increasingly focused on restoring health to sagebrush rangelands. Management is increasingly done in a proactive manner. For example, seeding controlled burns to prevent the establishment of nonnative plants, setting back succession in sagebrush stands to create a mosaic of sagebrush cover classes across the landscape, and adjusting grazing practices to retain tall grasses for nesting cover. Habitat management also involves restorative treatments designed to remove cheatgrass and other invasive/noxious weeds, removal of pinyon-juniper stands, and restoration of native species.

Several treatment types are used to manipulate sagebrush communities. Connelly et al. (2005:7-46 to 7-50) describes the mechanical, chemical, and biological techniques available and discusses their successes and challenges.

Given the current climate of vegetation management (i.e. restore/maintain plant/wildlife community health), vegetation management is not likely to be an important negative impact to sage-grouse populations in the Resource Area. As discussed in an earlier section of this Plan, several habitat management projects have been implemented, or proposed, that are designed to improve sage-grouse habitat. Further, the Utah Partners for Conservation and Development (UPCD), a collection of resource management agencies, NGO, and private individuals, recently established a Regional Team in the Resource Area. The purpose of the UPCD Regional Team is to increase communication, coordination, and sharing of resources and information with regards to habitat and watershed improvements in the Resource Area. Increased focus and coordination is likely to improve project planning, implementation, and outcomes.

Ongoing research is providing useful information about how sage-grouse use areas where vegetation treatments have been conducted (Robinson, unpublished data; Appendix B). Many acres have been treated in the Resource Area with the intent of improving sage-grouse habitat. Overall, we feel there is a great need to better monitor vegetation treatments in the Resource Area in an effort to expand our understanding of the influence of vegetation management on sage-grouse populations and habitats in the Resource Area.

## V. Conservation Strategy

One of the main purposes of this Plan is to provide a framework of strategies and associated actions that can be implemented to abate threats, address information gaps, and guide monitoring efforts. Strategies and actions listed below (the order is irrelevant) were developed by WDARM partners. Several other documents and publications provide recommendations and guidelines for management of sage-grouse populations and their habitats, many of which were reviewed in the Introduction of this Plan. Strategies developed by WDARM are designed to be specific to the local area while taking into consideration the guidelines provided at a rangewide level.

Implementation of strategies and actions is strictly voluntary on the part of WDARM partners. Despite this, we have designated for each strategy the public and private partners who might be involved in implementation. Designation does not imply responsibility or commitment of resources of any sort to implementing, initiating, or completing any actions. However, it does provide a framework of resources and expertise.

To help prioritize implementation of the Plan, efficiently use resources, and develop and secure funding, we have also identified which threats each strategy addresses. In addition, we list the aspects of sage-grouse ecology likely to be impacted by each strategy.

### A. Strategies and Actions

1. **Strategy:** Maintain and increase coordination and communication with agency and private partners.
  - 1.1. **Action:** Participate with and coordinate with the Central Region UPCD, Tooele County Natural Resource Group, Deep Creek Watershed partnership, Goshute Tribe, Tooele and Juab County Commissioners, SCDs, UFBF, and any other groups, as necessary.
  - 1.2. **Action:** Hold annual field tours to review projects, evaluate on-the-ground progress on the Plan, and share ideas.
  - 1.3. **Action:** Develop educational material appropriate for a broad recreationist audience to develop sensitivity to issues identified in the Plan.

**Partners:** UDWR, NRCS, BLM, USFS, UPCD, Goshute Tribe, County Commission, UFBF, USU Extension, County Weed Board, private partners  
**Threats Addressed:** Vegetation management, fire, OHV recreation, hunting  
**Aspects of Sage-grouse Ecology Addressed:** Seasonal habitat quality, population size, population distribution
2. **Strategy:** By 2010, reduce pinyon/juniper stands from sage-grouse use areas.
  - 2.1. **Action:** Remove pinyon/juniper trees from priority areas where action is warranted.
  - 2.2. **Action:** Revisit and retreat pinyon/juniper removal sites, as needed.

**Partners:** BLM, UDWR, USFS, NRCS, UFBF, private partners  
**Threats Addressed:** Vegetation management, fire, power lines, fences, and other tall structures, pinyon/juniper encroachment  
**Aspects of Sage-grouse Ecology Addressed:** Seasonal habitat quality, connectivity of seasonal habitat types

3. **Strategy:** By 2016, increase brood-rearing habitat quality in the Resource Area.
  - 3.1. **Action:** Work with the NRCS and private partners to develop projects that would increase brood-rearing habitat quality in the Resource Area.
  - 3.2. **Action:** Work with agency partners to develop projects that would increase brood-rearing habitat quality in the Resource Area.
  - 3.3. **Action:** Work with private and public partners to monitor effects of habitat improvement projects on vegetation and sage-grouse habitat use.
  - 3.4. **Action:** Where appropriate, reduce sagebrush canopy cover with mechanical or chemical treatments and reseed with ecologically appropriate seed mixes.

**Partners:** NRCS, BLM, USFS, UDWR, USU, UPCD, UFBB, private partners  
**Threats Addressed:** Vegetation management, fire, altered water distribution  
**Aspects of Sage-grouse Ecology Addressed:** Population size, population distribution, brood-rearing habitat quality, connectivity of seasonal habitat types
4. **Strategy:** Through 2016, maintain and protect winter habitat distribution and quality in the Resource Area.
  - 4.1. **Action:** Promote protection of winter habitat from fire.
  - 4.2. **Action:** Promote protection of winter habitat from OHV trail development and activities.
  - 4.3. **Action:** Update maps of crucial winter habitat areas and monitor winter habitat use areas for presence of sage-grouse.
  - 4.4. **Action:** In the event of fire, aggressively rehabilitate sites to prevent domination of invasive/noxious weed communities.

**Partners:** UDWR, BLM, USFS, NRCS, UFBB, UPCD, USU, private partners  
**Threats Addressed:** OHV recreation, vegetation management, fire, invasive/noxious weeds, development, power lines, fences, and other tall structures, renewable and nonrenewable energy development, pinyon/juniper encroachment  
**Aspects of Sage-grouse Ecology Addressed:** Winter habitat quality, connectivity of seasonal habitat types
5. **Strategy:** Reduce the threat of conversion of sagebrush stands to invasive/noxious weed communities.
  - 5.1. **Action:** Seed green-strips and/or fire breaks in crucial areas (to be identified).
  - 5.2. **Action:** Identify areas where fire suppression should be promoted to protect crucial habitat.
  - 5.3. **Action:** Maintain and/or increase fuels reduction projects in crucial areas (to be identified).
  - 5.4. **Action:** Work with agency and private partners to conduct vegetation treatments that restore functional plant groups to sagebrush communities.
  - 5.5. **Action:** Coordinate with noxious/invasive weed Coordinated Weed Management Area (CWMA) personal.

**Partners:** UDWR, BLM, County Weed Boards, USFS, Tooele County, Juab County, UPCD, USU, private partners  
**Threats Addressed:** Fire, invasive/noxious weeds, vegetation management, pinyon/juniper encroachment  
**Aspects of Sage-grouse Ecology Addressed:** Seasonal habitat quality, connectivity of seasonal habitat types

6. **Strategy:** Minimize the impact of excessive predation.
  - 6.1. **Action:** Modify power lines and wood fence posts (to remove raptor perches) in important sage-grouse areas, where feasible and where predator concerns have been identified.
  - 6.2. **Action:** Remove trees, remove/modify raptor perches, and maintain quality sagebrush habitat, where predation concerns on sage-grouse have been identified.
  - 6.3. **Action:** Maintain or increase site-specific predation management to consider all predator species (especially common ravens and red fox) where necessary and appropriate.
  - 6.4. **Action:** Initiate research on direct and indirect impacts of predation during each sage-grouse life history phase.
  - 6.5. **Action:** Coordinate management and research with USDA-WS.  
**Partners:** USDA Wildlife Services, UDWR, USU, private partners  
**Threats Addressed:** Predation; power lines, fences, and other tall structures, pinyon/juniper encroachment  
**Aspects of Sage-grouse Ecology Addressed:** Seasonal habitat quality, population size
  
7. **Strategy:** Work with public and private partners to implement livestock management plans that address seasonal needs of sage-grouse and livestock operations.
  - 7.1. **Action:** Incorporate appropriate livestock management in vegetation/habitat treatment projects.
  - 7.2. **Action:** Initiate research on the direct and indirect effects of livestock grazing on various aspects of sage-grouse life history.
  - 7.3. **Action:** Work with public and private partners to evaluate livestock management in crucial sage-grouse use areas.  
**Partners:** USU Extension, UFBB, BLM, USFS, private partners  
**Threats Addressed:** Livestock grazing, vegetation treatments  
**Aspects of Sage-grouse Ecology Addressed:** Seasonal habitat quality, connectivity of seasonal habitat types
  
8. **Strategy:** By 2016, increase population and habitat monitoring efforts in the Resource Area.
  - 8.1. **Action:** Encourage public and private partners to use techniques from Connelly et al. (2003) "Monitoring of Greater Sage-grouse Habitats and Populations."
  - 8.2. **Action:** In 2007, UDWR biologists will coordinate with Goshute Tribe biologists to identify sage-grouse lek sites and count birds on Tribal lands.
  - 8.3. **Action:** UDWR to enlist and coordinate private volunteers and/or other agency biologists to search for new leks and conduct lek counts on active leks.
  - 8.4. **Action:** Through 2016, test dead sage-grouse for West Nile Virus and any other parasites/pathogens of importance.
  - 8.5. **Action:** Secure funding to support additional research and monitoring on issue as identified in the Plan.
  - 8.6. **Action:** Increase outreach with private landowners to facilitate greater communication about sage-grouse distribution, ecology, and management.  
**Partners:** USU, USU Extension, BLM, USFS, UDWR, private partners  
**Threats Addressed:** Vegetation management, recreation, disease/parasites  
**Aspects of Sage-grouse Ecology Addressed:** Population size, population distribution, connectivity of populations/subpopulations

9. **Strategy:** Encourage use of this Plan in local, county, state, and federal natural resources planning efforts.
- 9.1. **Action:** Provide the Plan to all appropriate local, county, state, and federal natural resource agencies, departments, and personal.
- 9.2. **Action:** Review local, county, state, and federal plans and projects with the potential to impact sage-grouse and/or sagebrush habitats in the Resource Area.
- 9.3. **Action:** Participate in local, county, state, and federal natural resource planning efforts, committees, and working groups.
- Partners:** All
- Threats Addressed:** All
- Aspects of Sage-grouse Ecology Addressed:** All
10. **Strategy:** Minimize impacts of oil and gas development on sage-grouse and their habitat.
- 10.1. **Action:** Coordinate and communicate with BLM and USFS to ensure that adequate information/data is available for decision making process.
- 10.2. **Action:** Support recommendations that provide for temporal avoidance, minimization of tall structures, and avoid crucial habitat or use areas, where possible.
- 10.3. **Action:** Reduce fragmentation of sage-grouse habitat by oil and gas development activities.
- 10.4. **Action:** Minimize disturbance to sage-grouse associated with oil and gas development.
- 10.5. **Action:** Reduce cumulative impacts of oil and gas development.
- 10.6. **Action:** Share sage-grouse data with industry and encourage planning to reduce and/or mitigate for impacts.
- Partners:** BLM, USFS, UDWR, Tooele and Juab County Commissions, private partners
- Threats Addressed:** Renewable and nonrenewable energy development, power lines, fences, and other tall structures, roads, vegetation management, predation
- Aspects of Sage-grouse Ecology Addressed:** Seasonal habitat quality, connectivity of seasonal habitat types, connectivity of populations/subpopulations, population size, population distribution
11. **Strategy:** Minimize the amount of quality sage-grouse habitat eliminated by residential and commercial land development consistent with private property rights.
- 11.1. **Action:** Participate with County land-use decision makers in identifying key sage-grouse habitats.
- 11.2. **Action:** Maintain sagebrush environments of sufficient size and shape around developments in sage-grouse habitat.
- 11.3. **Action:** Encourage the voluntary use of conservation easements and other land protection vehicles with willing sellers in sage-grouse habitats.
- 11.4. **Action:** Educate rural residents about the importance of good grazing management in keeping small tracts weed free and capable of providing wildlife habitat.
- 11.5. **Action:** Work with public and private partners to maintain rural economies and viable ranching and agricultural enterprises.
- Partners:** Tooele and Juab County Commissions, UDWR, UFBF, NRCS, BLM, USFS, USU Extension, UACD, SCD, private partners
- Threats Addressed:** Livestock grazing, development, invasive/noxious weeds, vegetation

management

**Aspects of Sage-grouse Ecology Addressed:** Seasonal habitat quality, connectivity of seasonal habitats, population distribution

12. **Strategy:** By 2016, maintain or increase distribution and quality of mesic sites available to sage-grouse during summer months.
  - 12.1. **Action:** Work with public and private partners to develop mesic sites for sage-grouse associated with existing or new water developments.
  - 12.2. **Action:** Develop project planning tools (both printed material and on-the-ground examples) to illustrate successful, wildlife-friendly water developments.

**Partners:** NRCS, BLM, USFS, UDWR, UFBF, SCD, UACD, UPCD, private partners  
**Threats Addressed:** Altered water distribution  
**Aspects of Sage-grouse Ecology Addressed:** Breeding habitat quality, summer/brood-rearing habitat quality, population distribution
13. **Strategy:** Maintain or improve breeding habitat quality in the Resource Area.
  - 13.1. **Action:** Where appropriate, conduct vegetation manipulation to maintain open areas on lek sites.
  - 13.2. **Action:** Work with public and private partners to maintain nesting cover in crucial breeding areas.
  - 13.3. **Action:** Work with public and private partners to minimize disturbance to crucial areas during lek and nesting seasons.

**Partners:** NRCS, BLM, USFS, UDWR, UFBF, SCD, UACD, UPCD, private partners  
**Threats Addressed:** Livestock grazing, fire, vegetation treatment, invasive/noxious weeds, recreation, development, roads, power lines, fences, and other tall structures, pinyon/juniper encroachment  
**Aspects of Sage-grouse Ecology Addressed:** Breeding habitat quality, population distribution, population size
14. **Strategy:** Minimize the negative impacts of recreation on sage-grouse populations and their habitats.
  - 14.1. **Action:** Work with local, county, state, and federal planners and managers to minimize impacts of OHV trails and undeveloped roads on crucial sage-grouse habitat.
  - 14.2. **Action:** Work with law enforcement agencies to enforce existing and new laws, ordinances, and regulations specific to hunting/poaching, OHV recreation, and trespassing.
  - 14.3. **Action:** Work with OHV recreation groups to develop greater sensitivity and awareness to issues identified in this Plan.
  - 14.4. **Action:** If appropriate, work with public and private partners to restrict lek viewing opportunities during crucial time-periods and in crucial areas.
  - 14.5. **Action:** In a GIS system, evaluate where existing and proposed trails intersect crucial sage-grouse habitat.

**Partners:** Tooele and Juab County Commissions, BLM, USFS, UDWR, private partners.  
**Threats Addressed:** Recreation, development, hunting, roads  
**Aspects of Sage-grouse Ecology Addressed:** Seasonal habitat quality, population size, population distribution, connectivity of seasonal habitat types

## **B. Priority Evaluation**

In order to help prioritize strategies, actions, and most effectively allocate resources, we have assigned a rank of 'low', 'medium', 'high', or 'very high' to each threat with regards to its contribution to reduction in population health or habitat condition (Table 6). Again, given the stipulations regarding a lack of empirical, locally-based information in many cases, these rankings are based on the best information available to us and our implicit, experiential knowledge of the Resource Area. Ranking definitions are based on The Nature Conservancy's Conservation Action Planning process (TNC 2005). Rankings are provided to help highlight potential priorities for subsequent strategies and actions.

WDARM partners and others can use the rankings in Table 6, combined with the strategies and actions listed above, to prioritize implementation and direct resources to efficiently and effectively abate threats, and maintain and improve sage-grouse populations and their habitats in the Resource Area.

Table 6. Relative importance/contribution of individual threats to reducing or degrading aspects of sage-grouse populations in the WDARM Resource Area. Threats are described in the “Threat Analysis” section of this Plan. Rankings are as follows: L = low; M = medium; H = high; and VH = very high. Ranks are defined according to TNC (2005).

WDARM							
Threat	Reduced Population Size	Reduced Population Distribution	Reduced Breeding Habitat Quality	Reduced Late Summer/Fall Habitat Quality	Reduced Winter Habitat Quality	Reduced Connectivity of Seasonal Habitat Types	Reduced Connectivity of Populations & Sub-populations
Altered Water Distribution	-	VH	VH	H	L	L	H
Drought and Weather	M	H	M	M	L	L	-
Existing and New Fences	-	M	M	M	-	M	-
Home and Cabin Development	-	M	M	M	M	M	M
Power lines and Other Tall Structures	-	M	M	M	-	M	-
Renewable and Non-renewable Energy Development	-	M	M	M	-	L	L
Roads	-	M	M	M	M	M	M
Incompatible Vegetation Management	H	M	H	L	M	M	M
Poaching	H	L	-	-	-	-	-
Fire	-	-	VH	VH	VH	H	M
Incompatible Livestock Grazing	-	-	H	H	L	L	L
Recreation	VH	VH	H	M	VH	M	M
Invasive/Noxious Weeds	-	-	VH	VH	H	H	M
Parasites and Disease	M	M	-	-	-	-	-
Predation	VH	M	-	-	-	-	-
Pinyon-Juniper Encroachment	-	-	H	H	H	H	-
Conversion to Agriculture	-	-	L	L	-	-	-

#### IV. Literature Cited

- Aldridge, C. L. 1998. Status of the sage grouse (*Centrocercus urophasianus urophasianus*) in Alberta. Alberta Environmental Protection, Wildlife Management Division, and Alberta Conservation Association. Wildlife Status Report 13. Edmonton, Alberta, Canada.
- Anderson, D. R. 2001. The need to get the basics right in wildlife field studies. *Wildlife Society Bulletin* 29: 1294–1297.
- Apa, A. D. 1998. Habitat use and movements of sympatric sage and Columbian sharp-tailed grouse in southeastern Idaho. Dissertation, University of Idaho, Moscow, Idaho, USA.
- Armentrout, D. J., F. Hall, M. Dobel, and the Northeastern California Sage-grouse Working Group. 2004. Draft conservation strategy for sage-grouse (*Centrocercus urophasianus*) within the Buffalo-Skedaddle population management unit. Bureau of Land Management, Eagle Lake Office, Susanville, California, USA.
- Autenrieth, R. E. 1981. Sage grouse management in Idaho. Idaho Department of Fish and Game Wildlife Bulletin Number 9. Boise, Idaho, USA.
- Autenrieth, R. E., W. Molini, and C. E. Braun. 1982. Sage grouse management practices. Western States Sage Grouse Committee, Technical Bulletin 1, Twin Falls, Idaho, USA.
- Barnett, J. K., and J. A. Crawford. 1994. Pre-laying nutrition of Sage grouse hens in Oregon. *Journal of Range Management* 47: 114–118.
- Batterson, W. M., and W. B. Morse. 1948. Oregon Sage Grouse. Oregon Fauna Service, Oregon Game Commission, Portland, Oregon, USA.
- Beck, T. D. 1977. Sage grouse flock characteristics and habitat selection during winter. *Journal of Wildlife Management* 41: 18–26.
- Beck, T. D. I., and C. E. Braun. 1980. The strutting ground count: variation, traditionalism, management needs. *Proceedings of the Annual Conference of the Western Association of Fish and Wildlife Agencies* 60: 558–566.
- Beck, J. L., and D. L. Mitchell. 2000. Influences of livestock grazing on Sage Grouse habitat. *Wildlife Society Bulletin* 28: 993–1002.
- Beck, J. L., D. L. Mitchell, and B. D. Maxfield. 2003. Changes in the distribution and status of sage-grouse in Utah. *Western North American Naturalist* 63: 203–214.
- Bergerud, A. T. 1988. Population ecology of North American grouse. Pages 578–648 in A. T. Bergerud and M. W. Gratson, editors. *Adaptive strategies and population ecology of northern grouse*. University of Minnesota Press, Minneapolis, Minnesota, USA.
- Blanthorn, O. 1998. *A History of Tooele County*. Tooele, Utah.

- Blejwas, K. M., B. N. Sacks, M. M. Jaeger, and D. R. McCullough. 2002. The effectiveness of selective removal of breeding coyotes in reducing sheep predation. *Journal of Wildlife Management* 66: 451–462.
- Boyce, M. S. 1990. The red queen visits sage grouse leks. *American Zoologist* 30: 263-270.
- Braun, C. E. 1995. Distribution and status of Sage Grouse in Colorado. *Prairie Naturalist* 27: 1–9.
- Braun, C. E., T. Britt, and R. O. Wallestad. 1977. Guidelines for maintenance of Sage Grouse habitats. *Wildlife Society Bulletin* 5: 99–106.
- Bunnell, K. D. and J. T. Flinders. 1999. Restoration of sage grouse in Strawberry Valley, Utah 1998–99 report. Unpublished report to: Utah Reclamation Mitigation and Conservation Commission. Brigham Young University, Provo, Utah, USA.
- Bunnell, K. D., D. J. Bambrough, and J. T. Flinders. 2000. Revised progress report: Strawberry Valley Sage grouse recovery project. Brigham Young University, Provo, Utah, USA.
- Bunting, S. C., B. M. Kilgore, and C. L. Bushey. 1987. Guidelines for prescribed burning sagebrush-grass rangelands in the northern Great Basin. General Technical Report INT-231. **Publisher, City, state**, USA.
- Bureau of Land Management (BLM). 2005a. National Sage-grouse habitat conservation strategy. U.S. Department of the Interior, Bureau of Land Management, Washington, D.C., USA.
- Bureau of Land Management (BLM). 2005b. Salt Lake Fire Management Plan Environmental Assessment. UT-020-2004-0091. Salt Lake District, Salt Lake City, Utah, USA.
- Burkhardt, J. W., and E. W. Tisdale. 1969. Nature and successional status of western juniper vegetation in Idaho. *Journal of Range Management* 22: 264-270.
- Cannon, R. W., and F. L. Knopf. 1981. Lek numbers as a trend index to prairie grouse populations. *Journal of Wildlife Management* 45: 776–778.
- Churcher, C. S. 1959. The specific status of the New World red fox. *Journal of Mammalogy* 40: 513–520.
- Churcher, C. S. 1973. Red fox. Canadian Wildlife Service Hinterland Who's Who, Cat.CW69-4/5. **City**, Canada.
- Clary, W. P., and B. F. Webster. 1989. Managing grazing of riparian areas in the Intermountain Region. Gen. Tech. Rep. INT-263. U.S. Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, USA.
- Clary, W. P., N. L. Shaw, J. G. Dudley, V. A. Saab, J. W. Kinney, and L. C. Smithman. 1996. Response of a depleted sagebrush steppe riparian system to grazing control and woody plantings. USDA Forest Service Research Paper INT-RP-492. **City, State**, USA.

- Clary, W. P., and W. Leninger. 2000. Stubble height as a tool for management of riparian areas. Invited paper. *Journal of Range Management*. 53: 562–573
- Cline, N. L. 2006. Tooele County Population Estimates/Projections. Tooele County Planning and Economic Development Division, URL: <<http://www.tooeleconomicdevelopment.com/popest06.pdf>> Tooele, Utah, USA. Accessed August 21, 2006.
- Coggins, K. A. 1998. Sage grouse habitat use during the breeding season on Hart Mountain National Antelope Refuge. Thesis, Oregon State University, Corvallis, Oregon, USA.
- Connelly, J. W. 1982. An ecological study of Sage grouse in southeastern Idaho. Dissertation, Washington State University, Pullman, Washington, USA.
- Connelly, J. W., and C. E. Braun. 1997. Long-term changes in Sage grouse (*Centrocercus urophasianus*) populations in western North America. *Wildlife Biology* 3: 229–234.
- Connelly, J. W., W. J. Arthur, and O. D. Markham. 1981. Sage grouse leks on recently disturbed sites. *Journal of Range Management*. 34: 153–154.
- Connelly, J. W., H. W. Browsers, and R. J. Gates. 1988. Seasonal movements of Sage grouse in southeastern Idaho. *Journal of Wildlife Management* 52: 116–122.
- Connelly, J. W., W. L. Wakkinen, A. D. Apa, and K. P. Reese. 1991. Sage grouse use of nest sites in southeastern Idaho. *Journal of Wildlife Management* 55: 521–524.
- Connelly, J. W., R. A. Fischer, A. D. Apa, K. P. Reese, and W. L. Wakkinen. 1993. Renesting of Sage grouse in southeastern Idaho. *Condor* 95: 1041–1043.
- Connelly, J. W., K. P. Reese, and W. L. Wakkinen, M. D. Robertson, and R. A. Fischer. 1994. Sage grouse ecology report. Idaho Department of Fish and Game Job completion Report. W-160-R-19. Subproject 9. Boise, Idaho, USA.
- Connelly, J. W., M. W. Gratson, and K. P. Reese. 1998. Sharp-tailed grouse. The birds of North America, No. 354. The birds of North America, Inc., Philadelphia, Pennsylvania, USA.
- Connelly, J. W., M. A. Schroeder, A. R. Sands, and C. E. Braun. 2000. Guidelines to manage sage grouse populations and their habitats. *Wildlife Society Bulletin* 28: 967–985.
- Connelly, J. W., K. P. Reese, E. O. Garton, and M. L. Commons-Kemner. 2003a. Response of greater sage-grouse *Centrocercus urophasianus* populations to different levels of exploitation in Idaho, USA. *Wildlife Biology* 9: 335–340.
- Connelly, J. W., K. P. Reese, and M. A. Schroeder. 2003b. Monitoring of greater sage-grouse habitats and populations. University of Idaho, College of Natural Resources Experiment Station Bulletin 80, Moscow, Idaho, USA.

- Connelly, J. W., S. T. Knick, M. A. Schroeder, and S. J. Stiver. 2004. Conservation Assessment for Greater Sage-grouse and Sagebrush Habitats. Western Association of Fish and Wildlife Agencies. Unpublished Report. Cheyenne, Wyoming, USA.
- Connolly, G. E., and W. M. Longhurst. 1975. The effects of control on coyote populations: a simulation model. Division of Agricultural Sciences, University of California Bulletin 1872: 1–37.
- Crompton, B., R. Thacker, and T. Wright. 2006. Beer level answers for champagne level questions: quantifying the impacts of coalbed methane development on mule deer (*Odocoileus hemionus*) and greater sage-grouse (*Centrocercus urophasianus*). Utah Chapter of the Wildlife Society Annual Meeting, Moab, Utah, USA.
- Dalke, P. D., D. B. Pyrah, D. C. Stanton, J. E. Crawford, and E. F. Schlatterer. 1963. Ecology, productivity, and management of sage grouse in Idaho. *Journal of Wildlife Management* 27: 810–841.
- Danvir, R. E. 2002. Sage Grouse ecology and management in northern Utah sagebrush-steppe. A Deseret Land and Livestock Wildlife Research Report. Deseret Land and Livestock Ranch and the Utah Foundation for Quality Resource Management, Woodruff, Utah, USA.
- DeLong, A. K., J. A. Crawford, and D. C. DeLong, Jr. 1995. Relationships between vegetational structure and predation of artificial Sage grouse nests. *Journal of Wildlife Management* 59: 88–92.
- Drut, M. S., J. A. Crawford, and M. A. Gregg. 1994a. Brood habitat use by Sage Grouse in Oregon. *Great Basin Naturalist* 54: 170–176.
- Drut, M. S., W. H. Pyle, and J. A. Crawford. 1994b. Technical note: diets and food selection of Sage grouse chicks in Oregon. *Journal of Range Management* 47: 90–93.
- Dunkle, S. W. 1977. Swainson's Hawks on the Laramie plains, Wyoming. *Auk* 94: 65–71.
- Dunn, P. O., and C. E. Braun. 1986. Summer habitat use by adult female and juvenile Sage grouse. *Journal of Wildlife Management* 50: 228–235.
- Edelmann, F. B., M. J. Ulliman, M. J. Wisdom, K. P. Reese, and J. W. Connelly. 1998. Assessing habitat quality using population fitness parameters: a remote sensing/GIS-based habitat-explicit population model for Sage grouse (*Centrocercus urophasianus*). Technical Report 25. Idaho Forest, Wildlife and Range Experiment Station, Moscow, Idaho, USA.
- Ellis, K. L. 1984. Behavior of a lekking sage grouse in response to a perched golden eagle. *Western Birds* 15: 37–38.
- Eng, R. L. 1963. Observations on the breeding biology of male Sage Grouse. *Journal of Wildlife Management* 27: 841–846.

- Eng, R. L., and P. Schladweiler. 1972. Sage grouse winter movements and habitat use in central Montana. *Journal of Wildlife Management* 36: 141–146.
- Fischer, R. A. 1994. The effects of prescribed fire on the ecology of migratory Sage grouse in southeastern Idaho. Dissertation, University of Idaho, Moscow, USA.
- Fischer, R. A. 1996. An investigation on fire effects within xeric Sage grouse brood habitat. *Journal of Range Management* 49: 194–198.
- Fischer, R. A., A. D. Apa, W. L. Wakkinen, K. P. Reese, and J. W. Connelly. 1993. Nesting-area fidelity of Sage grouse in southeastern Idaho. *Condor* 95: 1038–1041.
- Flinders, J. T. 1999. Restoration of sage grouse in Strawberry Valley, Utah, 1998–99 report. Utah Reclamation Mitigation and Conservation Commission, Progress Report. Brigham Young University, Provo, Utah, USA.
- Friend, M., and J. C. Franson, editors. 1999. Field manual of wildlife diseases: general field procedures and diseases in birds. U.S. Geological Survey Biological Resources Division, Information and Technology Report 1999–2001. Madison, Wisconsin, USA.
- Gates, R. J. 1983. Sage grouse, lagomorph, and pronghorn use of a sagebrush grassland burn site on the Idaho National Engineering Laboratory. Thesis, Montana State University, Bozeman, Montana, USA.
- Gates, R. J. 1985. Observations of the formation of a Sage grouse lek. *Wilson Bulletin* 97: 219–211.
- Gese, E. M., R. L. Ruff, and R. L. Crabtree. 1996. Social and nutritional factors influencing the dispersal of resident coyotes. *Animal Behaviour* 52: 1025–1043.
- Gese, E. M., and R. L. Ruff. 1998. Howling by coyotes (*Canis latrans*): variation among social classes, seasons, and pack sizes. *Canadian Journal of Zoology* 76: 1037–1043.
- Gehrt, S. D., and W. R. Clark. 2003. Raccoons, coyotes, and reflections on the mesopredator release hypothesis. *Wildlife Society Bulletin* 31: 836–842.
- Gill, R. B. 1965. Distribution and abundance of a population of Sage grouse in North Park, Colorado. Thesis, Colorado State University, Fort Collins, Colorado, USA.
- Gilmore, R. M. 1946. Mammals in archeological collections from southwestern Pennsylvania. *Journal of Mammalogy* 27: 227–234
- Girard, G. L. 1937. Life history, habits, and food of the sage grouse, *Centrocerus urophasianus* Bonaparte. Publication 3. University of Wyoming, Laramie, Wyoming, USA.
- Godin, A. J. 1977. Wild mammals of New England. The Johns Hopkins University Press, Baltimore, Maryland, USA.

- Grahame, J. D., and T. D. Sisk, editors. 2002. Exotic and invasive species on the Colorado Plateau, <[http://cpluhna.nau.edu/Biota/invasive\\_exotics.htm](http://cpluhna.nau.edu/Biota/invasive_exotics.htm)> *In* Canyons, cultures and environmental change: an introduction to the land-use history of the Colorado Plateau. <<http://www.cpluhna.nau.edu/>> Accessed 16 March 2004.
- Gray, G. M. 1967. An ecological study of Sage grouse broods with reference to nesting, movements, food habits and sagebrush strip spraying in the Medicine Lodge drainage, Clark County, Idaho. Thesis, University of Idaho, Moscow, Idaho, USA.
- Gregg, M. A. 1991. Use and selection of nesting habitat by Sage grouse in Oregon. Thesis, Oregon State University, Corvallis, Oregon, USA.
- Gregg, M. A., J. A. Crawford, M. S. Drut, and A. K. DeLong. 1994. Vegetational cover and predation of sage grouse nests in Oregon. *Journal of Wildlife Management* 58: 162–166.
- Griner, L. A. 1939. A study of Sage grouse (*Centrocercus urophasianus*), with special reference to life history, habitat requirements, and numbers and distribution. Thesis, Utah State Agricultural College, Logan, Utah, USA.
- Gunnison Sage-grouse Rangewide Steering Committee. 2005. Gunnison Sage-grouse Rangewide Conservation Plan. Colorado Division of Wildlife, Denver, Colorado, USA.
- Hanf, J. M., P. A. Schmidt, and E. B. Groschen. 1994. Sage grouse in the high desert of central Oregon: results of a study, 1988–1993. U.S. Department of Interior, Bureau of Land Management Series P-SG-01, Prineville, Oregon, USA.
- Hartzler, J. E. 1972. An analysis of Sage Grouse lek behavior. Unpublished Dissertation, University of Montana, Missoula, Montana, USA.
- Heath, B. J., R. Straw, S. H. Anderson, J. Lawson, and M. Holloran. 1998. Sage-Grouse productivity, survival, and seasonal habitat use among three ranches with different livestock grazing, predator control, and harvest management practices. Federal Aid Job Completion Report. Wyoming Game and Fish Department, Cheyenne, Wyoming, USA.
- Hepworth, W. G. 1962. Diagnosis of diseases in mammals and birds. P-R Project FW-3-R-8, Job IW, Wyoming Game and Fish Commission, Cheyenne, Wyoming, USA.
- Hockett, G. A. 2002. Livestock impacts on the herbaceous components of Sage Grouse habitat: a literature review. *Intermountain Journal of Sciences* 8: 105–114.
- Holloran, M. J. 1999. Sage Grouse (*Centrocercus urophasianus*) seasonal habitat use near Casper, Wyoming. Thesis, University of Wyoming, Laramie, Wyoming, USA.
- Holloran, M. J. 2005. Greater Sage-grouse (*Centrocercus urophasianus*) population response to natural gas field development in western Wyoming. Dissertation, University of Wyoming, Laramie, Wyoming, USA.

*West Desert Greater Sage-grouse Conservation Plan*

- Honess, R. G. 1982. Cestodes of Grouse. Pages 161–164 in E. T. Thorne, N. Kingston, W. R. Jolley, and R. C. Bergstrom, editors. *Diseases of Wildlife in Wyoming*, Second Edition. Wyoming Game and Fish Department, Cheyenne, Wyoming, USA.
- Honess, R. G., and G. Post. 1968. Sage grouse coccidiosis. Pages 4–22 in Part 1: History of an epizootic in Sage Grouse. University of Wyoming Agricultural Experiment Station, Science Monograph 14, Laramie, Wyoming, USA.
- Hulet, B. V. 1983. Selected responses of Sage Grouse to prescribed fire, predation, and grazing by domestic sheep in southeastern Idaho. Thesis, Brigham Young University, Provo, Utah, USA.
- Hupp, J. W., and C. E. Braun. 1989. Topographic distribution of Sage grouse foraging in winter. *Journal of Wildlife Management* 53: 823–829.
- Jenni, D. A., and J. E. Hartzler. 1978. Attendance at a sage grouse lek: implications for spring censuses. *Journal of Wildlife Management* 42: 46–52.
- Johnson, G. D., and M. S. Boyce. 1990. Feeding trials with insects in the diet of Sage Grouse chicks. *Journal of Wildlife Management* 54: 89–91.
- Jolley, W. R. 1982. Protozoa. Pages 107–154 in E. T. Thorne, N. Kingston, W. R. Jolley, and R. C. Bergstrom, editors. *Diseases of Wildlife in Wyoming*, Second Edition. Wyoming Game and Fish Department, Cheyenne, Wyoming, USA.
- Klebenow, D. A. 1969. Sage grouse nesting and brood habitat in Idaho. *Journal of Wildlife Management* 33: 649–662.
- Klebenow, D. A. 1982. Livestock grazing interactions with Sage Grouse. Pages 113–123 in J. M. Peek and P. D. Dalke, editors. *Wildlife-livestock relationships symposium: Proceedings 10*. University of Idaho, College of Forestry, Wildlife, and Range, Moscow, Idaho, USA.
- Klebenow, D. A. 1985. Habitat management for Sage Grouse in Nevada. *World Pheasant Association Journal* 10: 34–46.
- Klott, J. H., and F. G. Lindzey. 1989. Comparison of sage and sharp-tailed grouse leks in south central Wyoming. *Great Basin Naturalist* 49: 275–278.
- Kovalchik, B. L., and W. Elmore. 1992. Effects of cattle grazing systems on willow-dominated plant associations in central Oregon. *Proceedings of the Symposium on Ecology and Management of Riparian Shrub Communities*. USDA, USFS, Intermountain Research Station General Technical Report INT-289, Ogden, Utah, USA.
- Leopold, A. 1933. *Game Management*. Macmillan, New York City, New York, USA.
- Lincoln County Sage Grouse Technical Review Team. 2004. Lincoln County sage grouse conservation plan. Lincoln County Coordinated Resource Management Steering Committee and the Nevada Division of Wildlife Resources. May 20, 2004. City, Nevada,

USA.

- Martin, N. S. 1970. Sagebrush control related to habitat and Sage grouse occurrence. *Journal of Wildlife Management* 34: 313–320.
- Meyers, O. B. 1992. Sage grouse habitat enhancement: effects of sagebrush fertilization. Dissertation, Colorado State University, Fort Collins, Colorado, USA.
- Miller, R. F., and J. A. Rose. 1995. Historic expansion of *Juniperus occidentalis* (western juniper) in southeastern Oregon. *The Great Basin Naturalist* 55: 37–45.
- Miller, R. F., and L. L. Eddleman. 2000. Spatial and temporal changes of sage grouse habitat in the sagebrush biome. Oregon State University Agricultural Experiment Station Technical Bulletin 151. Corvallis, Oregon, USA.
- Miller, R. F., T. J. Svejcar, and J. A. Rose. 2000. Impacts of western juniper on plant community composition and structure. *Journal of Range Management* 53: 574–585.
- Morgan County. 2006. Morgan County Consolidated Plan. Wasatch Front Regional Council, Coalville, Utah, USA.
- Mosley, J. C., P. S. Cook, A. J. Griffis, and J. O’Laughlin. 1997. Guidelines for managing cattle grazing in riparian areas to protect water quality: review of research and best management practices. Idaho Forest, Wildlife and Range Policy Analysis Group, University of Idaho, Moscow, Idaho, USA.
- Negative Population Growth. 2006. State Population Facts: Utah. URL: <http://www.npg.org/states/ut.htm> Accessed on August 21, 2006.
- Nelson, O. C. 1955. A field study of Sage Grouse in southeastern Oregon with special reference to reproduction and survival. Thesis, Oregon State College, Corvallis, Oregon, USA.
- Northwest Colorado Greater Sage-Grouse Working Group (GSGWG). 2004. Draft Northwest Colorado Greater Sage-grouse conservation plan. Colorado Division of Wildlife, Denver, Colorado, USA.
- Oakleaf, R. J. 1971. The relationship of Sage Grouse to upland meadows in Nevada. Thesis, University of Nevada, Reno, Nevada, USA.
- Paige, C., and S. A. Ritter. 1999. Birds in a sagebrush sea: managing sagebrush habitats for bird communities. Partners in Flight Western Working Group, Boise, Idaho, USA.
- Patterson, R. L. 1952. The Sage grouse in Wyoming. Sage Books, Inc., Denver, Colorado, USA.
- Peterson, R. L., R. O. Standfield, E. H. McEwen, and A. C. Brooks. 1953. Early records of red and the gray fox in Ontario. *Journal of Mammalogy* 34: 126–127.
- Petersen, B. E. 1980. Breeding and nesting ecology of female sage grouse in North Park,

Colorado. Thesis, Colorado State University, Fort Collins, Colorado, USA.

- Peterson, M., and N. Silvy. 1996. Reproductive stages limiting productivity of the endangered Attwater's prairie chicken. *Conservation Biology* 10: 1264–1276
- Phillips, J. B. 1990. Lek behaviour in birds: do displaying males reduce nest predation? *Animal Behaviour* 39: 555–565.
- Pitt, W. C., F. F. Knowlton, and P. W. Box. 2001. A new approach to understanding canid populations using an individual-based computer model: Preliminary results. *Endangered Species Update* 18: 103–106.
- Presnall, C. C., and A. Wood. 1953. Coyote predation on sage grouse. *Journal of Mammalogy* 34: 127.
- Pyle, W. H., and J. A. Crawford. 1996. Availability of foods of Sage grouse chicks following prescribed fire in sagebrush-bitterbrush. *Journal of Range Management* 49: 320–324.
- Rasmussen, D. I., and L. A. Griner. 1938. Life history and management studies of the Sage Grouse in Utah, with special reference to nesting and feeding habits. *Transactions of the North American Wildlife Conference* 3: 852–864.
- Remington, T. E., and C. E. Braun. 1985. Sage grouse food selection in winter, North Park, Colorado. *Journal of Wildlife Management* 49: 1055–1061.
- Rhoads, S. N. 1903. *The mammals of Pennsylvania and New Jersey*. Privately published, Philadelphia, Pennsylvania, USA.
- Robel, R. J., J. N. Briggs, A. D. Dayton, and L. C. Hulbert. 1970. Relationships between visual obstruction measurements and weight of grassland vegetation. *Journal of Range Management* 23: 295–297.
- Robel, R. J., J. A. Harrison, C. A. Hagen, J. C. Pitman, and R. R. Reker. 2004. Effect of energy development and human activity on the use of sand sagebrush habitat by lesser prairie-chickens in southwestern Kansas. *Transactions of the 69<sup>th</sup> North American Wildlife and Natural Resources Conferences*, Spokane, Washington, USA.
- Robertson, M. D. 1991. Winter ecology of migratory Sage grouse and associated effects of prescribed fire in southeastern Idaho. Thesis, University of Idaho, Moscow, Idaho, USA.
- Rogers, G. E. 1964. Sage Grouse investigations in Colorado. Technical Publication No. 16. Colorado Game, Fish and Parks Department, Denver, Colorado, USA.
- Rowland, M. 2004. Effects of management practices on grassland birds: greater sage-grouse. Northern Prairie Wildlife Research Center, Jamestown, North Dakota, USA.
- Sargeant, A. B., S. H. Allen, and R. T. Eberhardt. 1984. Red fox predation on breeding ducks in midcontinent North America. *Wildlife Monographs* 89 : pages.

- Sargeant, A. B., S. H. Allen, and J. O. Hastings. 1987. Spatial relations between sympatric coyotes and red foxes in North Dakota. *Journal of Wildlife Management* 51: 285–293.
- Savage, D. E. 1969. Relation of Sage grouse to upland meadows in Nevada. Nevada Fish and Game Commission, Job Completion Report, Project W-39-R-9, Job 12. Reno, Nevada, USA.
- Schroeder, M. A. 1997. Unusually high reproductive effort by Sage grouse in a fragmented habitat in north-central Washington. *Condor* 99: 933–941.
- Schroeder, M. A., J. R. Young, and C. E. Braun. 1999. Sage grouse (*Centrocercus urophasianus*). Pages ??? in A. Poole and F. Gill, editors. *The Birds of North America*, No. 425. The Birds of North America, Inc., Philadelphia, Pennsylvania, USA.
- Schroeder, M. A., and R. K. Baydack. 2001. Predation and the management of prairie grouse. *Wildlife Society Bulletin* 29: 24–32.
- Shaw, N. L. 1992. Recruitment and growth of Pacific willow and sandbar willow seedlings in response to season and intensity of cattle grazing. Pages 130–137 in W. P. Clary, E. D. McArthur, D. Bedunah, and C. L. Wambolt, compilers. *Proceedings-Symposium on ecology and management of riparian shrub communities*. USDA Forest Service General Technical Report INT-289, city, state, USA.
- Slater, S. J. 2003. Sage-grouse (*Centrocercus urophasianus*) use of different-aged burns and the effects of coyote control in southwestern Wyoming. Thesis, University of Wyoming, Laramie, Wyoming, USA.
- Sveum C. M. 1995. Habitat selection by Sage Grouse hens during the breeding season in south-central Washington. Thesis, Oregon State University, Corvallis, Oregon, USA.
- Sveum, C. M., W. D. Edge, and J. A. Crawford. 1998. Nesting habitat selection by Sage grouse in south-central Washington. *Journal of Range Management* 51: 265–269.
- Swenson, J. E., C. A. Simmons, and C. D. Eustace. 1987. Decrease of Sage grouse *Centrocercus urophasianus* after ploughing sagebrush steppe. *Biological Conservation* 41: 125–132.
- The Nature Conservancy (TNC). 2005. Conservation Action Planning: User Manual V 4.b. The Nature Conservancy, Arlington, Virginia, USA.
- Till, J. A. 1992. Behavioral effects of removal of coyote pups from dens. *Proceedings of the Vertebrate Pest Conference* 15: 396-399.
- Tooele County. 2005. Tooele Valley Trails Map. Tooele County Engineering Department, Tooele, Utah, USA. <[http://www.co.tooele.ut.us/eng\\_tcm maps.htm](http://www.co.tooele.ut.us/eng_tcm maps.htm)> Accessed on August 22, 2006.
- Trueblood, R. W. 1954. The effect of grass reseeding in sagebrush lands on Sage grouse

- populations. Thesis, Utah State Agricultural College, Logan, Utah, USA.
- U.S. Forest Service. 2005. Uinta National Forest Fire Management Plan. Unpublished Report, USDA Forest Service, Uinta National Forest, Provo, Utah, USA.
- Utah Department of Health. 2006. 2006 Utah Weekly Update. Utah Department of Health, Office of Epidemiology, Salt Lake City, Utah, USA.  
<<http://health.utah.gov/epi/diseases/wnv/DiseaseStatus/Track.htm>> Accessed on August 22, 2006.
- Utah Division of Oil, Gas, and Mining (OGM). 2006. Drilling activity database.  
<<http://www.ogm.utah.gov/oilgas/STATISTICS/drilling/1SPUD.htm>>. Accessed **date**.
- Utah Division of Water Resources. 2006. Precipitation Data.  
<<http://www.water.utah.gov/DroughtConditions/>>. Accessed **date**.
- Utah Division of Wildlife Resources (UDWR). 2002. Strategic Management Plan for Sage-grouse. Utah Department of Natural Resources, Division of Wildlife Resources, Publication 02–20, Salt Lake City, Utah, USA.
- Utah Division of Wildlife Resources (UDWR). 2006. Utah Sensitive Species List. Utah Department of Natural Resources, Division of Wildlife Resources. Salt Lake City, Utah, USA.
- Voigt, D. R., and B. D. Earle. 1983. Avoidance of coyotes by red fox families. *Journal of Wildlife Management* 47: 852–857.
- Wakkinen, W. L. 1990. Nest site characteristics and spring-summer movements of migratory Sage grouse in southeastern Idaho. Thesis, University of Idaho, Moscow, Idaho, USA.
- Wakkinen, W. L., K. P. Reese, and J. W. Connelly. 1992. Sage grouse nest locations in relation to leks. *Journal of Wildlife Management* 56: 381–383.
- Wallestad, R. O. 1971. Summer movements and habitat use by Sage grouse broods in Summer movements and habitat use by Sage grouse broods in central Montana. *Journal of Wildlife Management* 35: 129–136.
- Wallestad, R. 1975. Male Sage grouse responses to sagebrush treatment. *Journal of Wildlife Management* 39: 482–484.
- Wallestad, R., and D. Pyrah. 1974. Movement and nesting of Sage grouse hens in central Montana. *Journal of Wildlife Management* 38: 630–633.
- Wallestad, R. O., and P. Schladwiller. 1974. Breeding season movements and habitat selection of male Sage grouse. *Journal of Wildlife Management* 38: 634–637.
- Welch, B. L., J. C. Pedersen, and R. L. Rodriguez. 1988. Selection of big sagebrush by Sage grouse. *Great Basin Naturalist* 48: 274–279.

- Williams, C. L., K. Blejwas, J. J. Johnston, and M. M. Jaeger. 2003. **Title needed**. *Wildlife Society Bulletin* 31: 925–932.
- Winward, A. H. 2000. Monitoring the Vegetation Resources in Riparian Areas. General Technical Report RMRS-GTR-47. United States Department of Agriculture-Forest Service. Rocky Mountain Research Station, Ogden, Utah, USA.
- Wisdom, M. J., R. S. Holthausen, B. C. Wales, C. D. Hargis, V. A. Saab, D. C. Lee, W. J. Hann, T. D. Rich, M. M. Rowland, W. J. Murphy, and M. R. Eames. 2000. Source habitats for terrestrial vertebrates of focus in the Interior Columbia Basin: broad-scale trends and management implications. U.S. Department of Agriculture Forest Service General Technical Report PNW-GTR-485. Vol. 1–3. Pacific Northwest Research Station, Portland, Oregon, USA.
- Young, J. A., and R. A. Evans. 1981. Demography and fire history of a western juniper stand. *Journal of Range Management* 34: 501-506.
- Young, J. R. 1994. The influence of sexual selection on phenotypic and genetic divergence among sage grouse populations. Dissertation, Purdue University, West Lafayette, Indiana, USA.
- Young, J. R., C. E. Braun, S. J. Oyler-McCance, J. W. Hupp, and T. W. Quinn. 2000. A new species of Sage grouse (Phasianidae: *Centrocercus*) from southwestern Colorado. *Wilson Bulletin* 112: 445–453.
- Zablan, M. A. 1993. Evaluation of Sage grouse banding program in North Park, Colorado. Thesis, Colorado State University, Fort Collins, Colorado, USA.

**Appendix A**

West Desert Adaptive Resource Management Local Working Group  
Standard Operating Procedures

**Appendix B**  
Sage-grouse Research Project Reporting